

Effect of Prey and Predator age on the Feeding Preference and rate of Predation by two Predators *Coccinella transversalis* Fab. and *Cheilomenes sexmaculatus* (Coleoptera : Coccinellidae)

R. VEERAVEL and P. BASKARAN

Department of Agricultural Entomology, Faculty of Agriculture
Annamalai University
Annamalainagar-608 002
Tamilnadu

ABSTRACT

The feeding preference of two coccinellid predators namely, *Coccinella transversalis* F. and *Cheilomenes sexmaculatus* F. on different instars and forms of the aphid, *Aphis gossypii* Glov. and predator age-dependent interference on the rate of predation were investigated. Significant differences were observed among the predatory stages for various prey ages. Older predators fed on older preys. However, adult beetles did not show any marked preference and fed on all stages and forms of preys. The various combinations of adults and grubs at different levels significantly affected the prey consumption levels. Among the various treatments, the treatment with 12 predators consumed the maximum number of aphids at all prey levels followed by 12 adults + one instar grub indicating the addition of a grub did not increase the prey consumption of adults.

KEY WORDS : *Cheilomenes sexmaculatus* , *Coccinella transversalis*, predation, *Aphis gossypii*

Coccinellids are mostly polyphagous beetles with a wide host range of accepted food ranging from coccids to aphids (Hodek, 1965) and the rate of feeding has been reported to be widely changing depending on the stage, and sex of the predator and also on the size of the prey. The prey consumption is known to be positively correlated with the population density of the prey (Popov, 1960) and the age of the predator has influence on the prey consumption (Choi and Kim, 1985). Likewise, aged grubs appeared to have a depressing effect on the feeding of younger stages (Murdoch and Andrew Sih, 1978). The present investigation was initiated to study the prey age preference of predators and predator age interference on the rate of predation under laboratory conditions.

MATERIALS AND METHODS

Potted brinjal plants of the var. Panruti local were maintained continuously in the insectary. *Aphis gossypii* Glover, collected from

brinjal fields were released on the third leaf of 40 days old plants. The plants with aphids were then covered with a muslin cloth bag on bamboo frame and the aphids were allowed to multiply. In another set of plants, *A. gossypii* were introduced and allowed to reproduce so as to have a thick population. The plants were then arranged in two rows within an iron cage covered with muslin cloth. Adult coccinellids of both *Coccinella transversalis* F. and *Cheilomenes sexmaculatus* F. collected from fields were then released separately. The beetles were allowed to feed on aphids and multiply. The plants were replaced with fresh ones as and when required so as to maintain a continuous supply of aphids for the growth and development of the coccinellids.

Individual predatory grubs of I, II, III and IV stage and adults from the stock culture were taken in Petri dishes separately and provided with 20 numbers of either I, II, III and IV instar aphid nymphs, parthenogenetic female or winged aphids on brinjal leaves. The experi-

ment was replicated three times and conducted at a temperature of $28 \pm 1^{\circ}\text{C}$. The preference for a particular category of prey by the predator was worked out as per the number consumed in 24 h time. A set of control dishes was also maintained to find out the natural mortality of the aphids. The mean level of prey preference shown by different predatory stages was tested using analysis of variance for two way classification and were compared by using DMRT.

To evaluate the relative contribution of different components of predation, three experiments were carried out as per the procedures described by Murdoch and Andrew Sih (1978). In each case, the presence or absence of a component constituted the treatment. The treatments were fixed mainly to study the functional and developmental responses of predators. The experiment was carried out in Petri dishes. The design was as follows:

- A - six adults (Functional response)
- B - Twelve adults (Functional response)
- C - One male + two females + four instar grub (Functional & Developmental response)
- D - One male + two females (Functional response)

- F - Four II instar grubs (Functional response)
- G - Four III instar grubs (Functional response)
- H - Four IV instar grubs (Functional response)
- I - Six adults + one I instar added/day (Functional & Developmental response)
- J - Twelve adults + one I instar added/day (Functional & Developmental response)
- K - One male + two females + four I instar grubs added/day (Functional & Developmental response)

Predator numbers were ensured each day. The density of adults was maintained by replacing those that died. However, young ones were allowed to develop and were not replaced when they died. Every day, aphids of constant size were added at levels of 50, 100, 150, 200, 250, 300, 350, 400, 450 and 500 as prey. Each morning the prey insects surviving from the previous day were removed and counted. Each experiment was replicated three times and control dishes were maintained with the same number of aphids but without predators.

Analysis of variance technique was used to analyse the data and the means were compared using DMRT.

Table 1. Number of prey consumed/day by the coccinellid predators* (Mean of three observations)

| Stage of the predator | Stage of the Aphid | | | | | |
|------------------------|--------------------|---------|---------|-------|------------------------|--------------|
| | I | II | III | IV | Parthenogenetic female | Winged aphid |
| <i>C.transversalis</i> | | | | | | |
| I stage grub | 7.6a | 2.0b | 2.3b | 2.0b | 0.0 | 0.0 |
| II stage grub | 1.7c | 4.3a | 4.3a | 4.0a | 2.7b | 0.0 |
| IV stage grub | 1.0c | 5.7a | 4.0a | 3.7b | 3.7b | 3.3b |
| IV stage grub | 2.3d | 7.3ab | 8.3a | 4.7c | 3.0d | 2.3d |
| Male | 5.7c | 12.0ab | 13.7a | 7.3c | 4.3c | 3.0d |
| Female | 10.7d | 12.0bcd | 14.0abc | 15.7a | 14.3ab | 1.33e |
| <i>M.sexmaculatus</i> | | | | | | |
| I stage grub | 6.0a | 2.0b | 1.0bc | 0.7c | 0.0 | 0.0 |
| II stage grub | 2.7b | 2.0b | 2.3ab | 3.0a | 2.3ab | 0.3c |
| III stage grub | 3.7ab | 4.7a | 3.7ab | 3.0ab | 4.0ab | 0.0 |
| IV stage grub | 1.7b | 2.0ab | 2.7a | 1.3b | 2.7a | 0.3c |
| Male | 10.7ab | 11.7a | 11.3ab | 4.3c | 1.0d | 0.0 |
| Female | 14.0ab | 15.0a | 11.7c | 9.0d | 5.7e | 1.3f |

* Means followed by similar letters within the predator species in vertical columns are not different statistically ($p=0.05$) by D.M.R.T.

Table 2. Age - dependent interference in *C. transversalis* (I) and *C. sexmaculatus* (II) in terms of prey aphid consumption at various combination of stages of predators* (Mean of three observations)

| Prey aphid density (No./leaf) | Predator | Number of prey consumed/day | | | | | | | | | | | | S.E |
|----------------------------------|----------|-----------------------------|-------|-------|-------|-------|-------|-------|--------|--------|-------|-------|------|--------|
| | | A | B | C | D | E | F | G | H | I | J | K | L | |
| 50 | I | 42ab | 50a | 35ab | 32b | 30b | 34a | 40ab | 38b | 43ab | 50a | 36ab | 5c | 6.870 |
| | II | 40c | 48b | 32e | 29f | 27g | 36d | 32e | 32e | 40c | 50a | 31e | 6h | 0.0213 |
| 100 | I | 84cd | 100a | 68g | 63h | 61h | 67g | 60e | 75f | 86c | 98ab | 74f | 12i | 0.883 |
| | II | 81c | 97a | 60g | 42i | 37j | 58h | 72e | 68f | 77d | 89b | 60g | 13k | 0.136 |
| 150 | I | 140abc | 150a | 105d | 100d | 84d | 94d | 101d | 110bcd | 118bcd | 148ab | 106d | 20e | 11.070 |
| | II | 132b | 140aa | 68g | 81de | 72f | 78e | 80e | 95c | 84d | 95c | 82d | 22h | 1.040 |
| 200 | I | 145d | 165b | 124g | 120gh | 115i | 130ef | 121gh | 132e | 160c | 180a | 118h | 25j | 1.350 |
| | II | 138a | 145a | 72e | 102c | 94cd | 98cd | 92d | 97cd | 125b | 120b | 96cd | 28f | 2.987 |
| 250 | I | 162c | 172a | 140f | 131gh | 122i | 133g | 145e | 141f | 152d | 167b | 138f | 30j | 1.430 |
| | II | 142b | 148a | 78h | 105g | 102g | 115e | 110f | 117de | 134c | 142b | 120d | 30i | 0.832 |
| 300 | I | 168c | 178a | 142g | 134i | 129j | 137h | 151e | 147f | 156d | 172b | 142g | 28k | 0.891 |
| | II | 150c | 155b | 84k | 112i | 110j | 120g | 117h | 121f | 142d | 168a | 129e | 31i | 0.105 |
| 350 | I | 170b | 185a | 148e | 137g | 132h | 142f | 156d | 152de | 161c | 182a | 145ef | 32i | 1.150 |
| | II | 152c | 161b | 91i | 118h | 115hi | 130fg | 132f | 128g | 145d | 172a | 134e | 35 | 0.653 |
| 400 | I | 172c | 190a | 150fg | 141i | 134j | 147gh | 162d | 158de | 169cd | 185b | 152f | 35k | 1.157 |
| | II | 161b | 169b | 98f | 125d | 121de | 135cd | 140c | 132cd | 155bc | 189a | 140c | 37g | 5.890 |
| 450 | I | 175c | 192a | 157e | 143g | 139h | 152f | 167d | 160e | 172cd | 190ab | 156e | 40i | 1.414 |
| | II | 168bc | 172b | 100e | 132cd | 129cd | 138cd | 145cd | 141cd | 158bc | 194a | 142cd | 41f | 1.568 |
| 500 | I | 182b | 200a | 161e | 148f | 142g | 159ef | 159d | 162e | 175c | 198a | 160e | 41h | 0.871 |
| | II | 171b | 175b | 110g | 139e | 135f | 140e | 157d | 149e | 162c | 197a | 151e | 43h | 1.860 |
| Mean | I | 144.0 | 158.2 | 123.0 | 11.49 | 108.8 | 119.5 | 129.2 | 127.5 | 139.2 | 157.0 | 122.7 | 26.8 | |
| | II | 133.5 | 141.0 | 79.3 | 98.5 | 94.5 | 104.8 | 107.7 | 108.0 | 122.2 | 141.6 | 108.5 | 28.6 | |

A = Six adults; B = Twelve adults; C = One male + two females + four I instar grubs; D = One male + two females; E = Four I instar grubs; F = Four II instar grubs; G = Four III instar grubs; H = Four IV instar grubs; I = Six adults + one I instar-day; J = Twelve adults + one I instar added; K = One male + two females + four I instar grubs; L = Control

* Means followed by similar letters in vertical columns are not different statistically ($p=0.05$) by D.M.R.T.

RESULTS AND DISCUSSION

Feeding preference shown by the various stages of predators for different instars of aphids are given in Table 1. Significant difference in preference for prey of various ages was evident among the predators. The first stage grubs killed preys of all instars of aphids and the total consumption was however lower compared to second stage grubs that killed and consumed the parthenogenetic females in addition. The third and fourth stage grubs consumed winged aphids and the adult predators consumed appreciable number of all the stages. Among the sexes, the female consumed more number of aphids than the male. The mean consumption of II, III and IV instar aphids by all the stages of the predator remained statisti-

cally on par in *C. transversalis*. Among the predatory grubs, the fourth instar and third instar consumed the maximum number of aphids in the case of *C. transversalis* and *C. sexmaculatus* respectively. As the grubs increased in age, they consumed older instars of aphids and the prey age preference shown by predatory grubs for the first instar aphids might be due to the fragile nature of these preys. As the grub increased in age and size, it attacked larger aphids with ease due to its increased mobility, activity and strength. Adult beetles fed on all age groups and forms of aphid, and no specific preference among the prey was seen. Saharia (1981) made similar observations in *Coccinella repanda* Feb., and no feeding

preference was shown by grown up grubs and adult predators for prey aphid groups.

The age dependent interference of the predators to prey consumption was determined by taking various combination of prey stages and allowing them to feed on known number of preys. The results presented in Table 2 showed that the various combination of adults and grubs at different levels significantly influenced the prey consumption. The treatment with 12 adult predators (B) resulted in maximum number of aphids eaten at all prey levels followed by the treatment with 12 adults + one first instar grub (J) and were statistically on par. The addition of one first instar grub in the combination did not increase the feeding rate of adults. Treatment A with six adults consumed appreciable number of aphids (144.0).

However, the juveniles showed variation between the different age groups viz., I, II, III and IV instars. The first instar grub, being very small in size consumed less number of aphids compared to other stages of grubs. Hence, when the grub developed into II, III and IV instars, the developmental response operated and the predator consumed more number of aphids. Among the grubs, the third instar grub consumed the maximum mean number of aphids (129.2). The prey consumption in treatments C and F and K were statistically on par. In *C.sex-maculatus*, the prey consumption by twelve adults + one first instar grub, (J) remained highest and statistically identical.

Since, both grub and adult coccinellids feed on the same kind of prey, possible existence of interference among them while feeding was expected but the results revealed no such interference since the feeding was not af-

fected either by the presence of grubs or adults in the feeding arena. Among the grubs, the third stage grubs consumed maximum number of aphids than the other stages. This increased feeding might be due to its increased age and capture efficiency by a process of learning as reported by Murdoch (1971) in *Notonecta hoffmanni* Hungerford. However, the fourth stage grubs consumed less number of aphids possibly due to entering prepupation. Another possible reason might be the increase in age of predatory grubs resulting in the rapidity of predator response following prey contact thereby taking less time to catch and consume more number of prey (Murdoch, 1971; Wratten, 1973).

REFERENCES

- CHOI, S.Y. and KIM, G.H. 1985. Aphidivorous activity of a coccinellid beetle, *Harmonia axyridis* Pallas (Coleoptera; Coccinellidae). *Korean Plant Prot.*, 24 11-14.
- HODEK, I. 1965. The predation of *Coccinella septempunctata* on *Aphis fabae* on sugarbeet. *Acta Ent. Bhenoslov.*, 62(4), 241-253.
- MURDOCH, W.W. 1971. The developmental response of predators to changes in prey density. *Ecology*, 52, 132-137.
- MURDOCH, W.W. and ANDREW SIH. 1978. Age-dependent Interference in a predatory insect. *J. Animal Ecol.*, 47, 581-592.
- POPOV, P. 1960. Role of *Coccinella septempunctata* L. in the biological control of aphids. *Selskistop Misul sofiya*, 152- 153.
- SAHARIA, D. 1981. Biology of *Coccinella repanda* Thunb. an important predator of mustard aphid *Lipaphis erysimi* (klth.) in Assam. *J. Res. Assam Agric. Univ.*, 2, 66-73.
- WRATTEN, S.D. 1973. The effectiveness of the coccinellid beetles *Adalia bipunctata* (L.) as a predator of the lime aphid *Eucaplipterus tibiae* L. *J. Animal Ecol.*, 42, 801.