DOI: *10.15613/fijrfn/2024/v11i2/47845*
ISSN (Print) : 2349-8897
***FoodSci*: Indian Journal of Research in Food Science and Nutrition,** *Vol 11(2), 69-77, July-December 2024* *ISSN (Online) : 2350-1006*

Acrylamide in Food: From Formation to Regulation and Emerging Solutions for Safer Consumption

S. Abhigna*, Mayuri Kulkarni, Nimisha Khandalkar and Aishwariya V. Patil

Department of Food Technology, M. S. Ramaiah University of Applied Sciences, Bengaluru -560054, India; abhigna2002@gmail.com

Abstract

Acrylamide is a heat-induced chemical that was first identified in food in 2002. It ismostlycreated when reducing sugars and asparagine undergo the Maillard reaction, which occurs during high-temperature cooking techniques like roasting, baking, and frying. Its synthesis is also aided by other processes, including lipid breakdown. Crispy bread, coffee, baked foods, cereals, and potato items are common sources. Acrylamide has caused serious health concerns due to its neurotoxic, genotoxic, and carcinogenic qualities as well as the International Agency for Research on Cancer's designation of it as a potential human carcinogen. Epidemiological research indicates that eating acrylamide increases the risk of cancer, especially in high-exposure populations. Reduced acrylamide levels are the goal of regulatory initiatives by organizations like the FDA and EFSA, yet difficulties still exist because detection techniques are expensive. Using antioxidants like polyphenols, changing cooking methods, and applying enzyme treatments are examples of current mitigating tactics. This review emphasizes the need for creative solutions to improve food safety by highlighting the mechanics of acrylamide generation, health hazards, dietary sources, and detection and mitigation strategies.

Keywords: Acrylamide, Carcinogenic Risks, Food Safety, High-Temperature Cooking

1. Introduction

Following the discovery in food in 2002, acrylamide- a chemical molecule produced during high-temperature cooking methods like frying, roasting, and baking- has attracted a lot of attention. The main cause of its creation is the Maillard reaction, which occurs when reducing carbohydrates like glucose and fructose and the amino acid asparagine are heated beyond 120°C[1]. Acrylamide can also form through lipid degradation processes that involve intermediates like acrolein in addition to the Maillard reaction[2]. Bread, coffee, cereal items, and snacks made from potatoes are foods that are frequently linked to acrylamide[3]. Because of its neurotoxic, genotoxic, and carcinogenic qualities, the chemical has caused health concerns ever since it was discovered in foods in 2002. According to the International Agency for Research (IARC) on Cancer, acrylamide is a likely human carcinogen. Epidemiological research has linked dietary acrylamide consumption to a higher risk of developing some types of cancer, especially in groups where exposure levels (Figure1) are high[4,5].

Guidelines have been put in place by regulatory agencies including the U.S. Food and Drug Administration (FDA) and the European Food Safety Authority (EFSA) to monitor and reduce the amount of acrylamide in food products. For example, EFSA published Regulation 2017/2158, which outlines doable steps to lower acrylamide in foods such as baked goods and products made from potatoes[6,7]. Despite these initiatives, the high expense of analytical methods such as liquid chromatography-mass spectrometry makes it difficult to precisely measure the levels of acrylamide in various food matrices[8].

Using enzymes like asparaginase to reduce precursors in food and altering cooking techniques, including lowering frying temperatures and times, are the main focuses of mitigation measures. Furthermore, it has been discovered that antioxidants, such as dietary polyphenols, mitigate the toxicity caused by acrylamide, providing encouraging opportunities for risk mitigation[8,9].

*Author for correspondence

Acrylamide in food is discussed in this review along with its production during high-temperature cooking, potential health hazards such as cancer and neurotoxicity, and detection and reduction techniques. Changing cooking techniques is one way to reduce acrylamide.

2. Formation of Acrylamide in Foods

When foods high in carbohydrates are cooked at high temperatures, acrylamide, a substance that is categorized as a possible carcinogen, is created. Let's see the various processes that led to its formation.

2.1 Pathway of the Maillard Reaction

One of the main processes to produce acrylamide is the Maillard reaction, which occurs particularly in meals high in reducing sugars and amino acids, like coffee, cereals, and potatoes. Acrylamide is one of the several intermediate compounds that are produced by this intricate, non-enzymatic browning reaction, which takes place at high temperatures (over 120°C)(Figure 2). Acrylamide is produced during this reaction when asparagine combines with reducing carbohydrates such as fructose and glucose. This is especially noticeable in dishes like roasted coffee beans or fried potatoes that contain a lot of carbohydrates. Although the process is essential for giving cooked foods their flavour and colour, it also produces toxic byproducts like acrylamide. According to studies, altering the cooking time and temperature might lessen the severity of the Maillard reaction and minimize the generation of acrylamide[11,12].

2.2 Decomposition of Asparagine

An amino acid called asparagine is essential for the synthesis of acrylamide. Acrylamide is a byproduct of asparagine's breakdown under heat, especially when it interacts with reducing sugars. In starchy foods like potatoes, where high asparagine levels lead to increased acrylamide formation, asparagine breakdown is essential to the Maillard process. Acrylamide generation can be considerably decreased by lowering asparagine concentrations in raw dietary ingredients before cooking, either by genetic selection or agricultural techniques. Pre-cooking techniques like blanching or soaking may also diminish asparagine levels and, consequently, acrylamide in the finished product, according to studies[3,14].

2.3 Interactions between Amino Acids and Lipids

Lipid and amino acid interactions, especially during frying, provide another avenue to produce acrylamide.



Figure 1. The burden of disease due to dietary exposure to acrylamide (Source: Mihalache, O. A., and Dall'Asta, C.; 2024. DOI: 10.1016/j.fct.2024.114699)[10].




Figure 2. Mechanism of acrylamide formation in food (Source: Huang, C., *et al*; 2021 DOI: 10.53365/nrfhh/142376)[13].

Frying's high temperatures cause lipids to break down, producing volatile chemicals that can react with amino acids—especially asparagine—to produce acrylamide. Fried snacks and other foods high in fat are more likely to have this pathway. When lipids and oils used in deep-frying combine with amino acids found in starchy foods like potatoes and cereals, research shows that the resulting chemicals can aid in the creation of acrylamide[15,16].

2.4 The Impact of the Food Matrix

Acrylamide production is significantly influenced by the food matrix, which consists of the food's structure and substance. Food's pH, moisture content, and starch content all have a big impact on how much acrylamide is formed. High-heat cooking techniques like frying and baking tend to increase the production of acrylamide in foods with a high starch-to-water ratio, like potatoes. Because water slows down the Maillard reaction and lessens the synthesis of acrylamide, the moisture content is crucial. Additionally, because the Maillard reaction proceeds more quickly in meals with a higher pH (alkaline foods), acrylamide generation may be enhanced[17,18].

2.5 Different Cooking Conditions

Temperature, cooking duration, and moisture content are some of the variables that affect the development of acrylamide during cooking. Acrylamide is produced because of the Maillard reaction between sugars and amino acids, especially asparagine, which is facilitated by higher temperatures, particularly at 120°C(Figure 3). Acrylamide levels are considerably raised when meals like potatoes are fried at temperatures higher than 170°C[18]. Furthermore, cooking time is important since longer cooking times can raise the amounts of acrylamide, while overcooking can cause it to break down into different components[19]. Because they cook in a drier environment, foods with lower moisture content—like baked goods or fried potatoes—tend to produce more acrylamide. On the other hand, acrylamide levels are typically reduced when foods with a high moisture content are boiled or steamed. Acrylamide formation is also significantly influenced by the type of diet and its composition, especially its starch content, with foods high in starch being more likely to produce it. Therefore, controlling the cooking process can assist prevent the development of acrylamide while preserving the quality of the meal[20].

3. Sources of Acrylamide in the Diet

Many foods, particularly those prepared at high temperatures using techniques like frying, roasting, and baking, contain acrylamide. The following are the primary foods that increase exposure to acrylamide.

3.1 Potato Products

Foods made from potatoes, especially those that are fried, roasted, or baked at high temperatures, are a major

source of acrylamide in the diet. One of the main sources of acrylamide is fried potato products, such as French fries and potato chips. When potatoes are cooked at temperatures higher than 120°C, the Maillard reaction takes place, resulting in the formation. Temperature and cooking time both raise acrylamide levels. Crispy or browned potatoes typically contain the most acrylamide[22,23].

3.2 Cereal-Based Foods

Acrylamide is frequently found in bread, biscuits, breakfast cereals, and cereal-based crackers. High temperatures during baking trigger the Maillard reaction, which results in the formation of acrylamide in the meal. Products that are roasted or darker tend to have higher quantities of acrylamide. The amount of acrylamide in various foods can vary depending on the kind of cereal and the baking circumstances[22].

3.3 Coffee

Roasted coffee beans are a significant source of acrylamide, with the levels varying based on the roasting process. During roasting, the beans are exposed to high temperatures above 200°C, which triggers the Maillard reaction, leading to the formation of acrylamide. Darker roasts, which undergo longer roasting times, typically contain higher levels of acrylamide compared to lighter roasts. This is due to the prolonged heat exposure, which intensifies the chemical reactions within the beans. As a result, regular consumption of coffee, especially dark roasts, can contribute to increased acrylamide intake, a concern for individuals mindful of its potential health risks[22,23].




Figure 3. Effects of power and time in MW roasting on acrylamide formation amount in salted and unsalted pistachio types (Source:Asadi *et al.*; 2020DOI: 10.1002/fsn3.1588)[21].

3.4 Baked Products

Baked goods such as cakes, cookies, pastries, and toasted bread are common sources of acrylamide in the diet. The Maillard reaction, which takes place at high temperatures typically above 120°C, is responsible for acrylamide formation during baking. Foods that are baked until golden brown or crispy tend to have higher levels of acrylamide. This is particularly evident in products that undergo longer baking times or are cooked at higher temperatures, as the heat promotes the reaction between sugars and amino acids like asparagine. Therefore, baked goods with a darker, crispier texture generally contain higher acrylamide levels, making them a significant dietary source of this compound[22].

3.5 Fried Foods

Fried foods, including deep-fried snacks and meats, represent another major source of acrylamide. The high temperatures involved in frying, often exceeding 170°C, accelerate the Maillard reaction, which results in acrylamide formation. Fried foods that are golden brown or have a crispy exterior are particularly high in acrylamide. This is because the longer foods are exposed to heat, and the more they brown or crisp up, the greater the formation of acrylamide. Therefore, it is advisable to fry foods at lower temperatures or for shorter durations to reduce acrylamide formation, although this may affect the texture and taste of the food[23].

3.6 Biscuits and Crackers

Another significant source of acrylamide in the diet is dry baked foods like biscuits, crackers, etc. High temperatures are usually used to bake these foods to give such a crunchy texture, which promotes the Maillard process and forms acrylamide. The amount of acrylamide in the product increases with its colour. Additionally, the usage of refined flour and sugars raises the levels of acrylamide[22,23].

3.7 Crispy Bread and Toasted Bread

Acrylamide can also be found in considerable quantities in crisp breads like rye and toasted bread (Figure 4).




Figure 4. Various foods contribute to Acrylamide in the diet. (Source:Barón Cortés, W. R., *et al.*; 2021 DOI: 10.1016/j.jfca.2021.103934)[24].

Bread is subjected to high temperatures during the toasting process, which triggers the Maillard reaction and promotes the production of acrylamide. Acrylamide levels are higher in darker-toasted bread than in lightly toasted bread[23].

4. Health Implications of Exposure to Acrylamide

According to national dietary patterns and food preparation methods, varying levels of acrylamide (AA) have been reported in foods across different countries. Potato-based products, coffee, and baked goods are identified as the primary sources. Direct exposure to AA typically occurs through the consumption of high-carbohydrate foods such as potato crisps, chips, roasted cereals, and bread. Additionally, indirect exposure to AA can result from food packaging containing polyacrylamide, which may leave residual monomers[25]. Acrylamide is commonly ingested through oral routes, such as dietary intake, drinking water, or smoking. Studies have shown that rats have demonstrated that bioavailability through diet ranges between 30% and 40%[26]. Once absorbed, acrylamide undergoes metabolism through the cytochrome P450 oxidase system, specifically CYP2E1, resulting in the formation of glycidamide (GA). Alternatively, it can directly react with glutathione to produce N-acetyl-S-(2-carbamoyl ethyl) cysteine (AAMA)[27].Exposure to this is associated with neurotoxicity, impaired fertility, cancer risk and so on.

4.1 Carcinogenicity

In 2015, 17.5 million cancer cases and 8.7 million deaths were reported globally, with 208.3 million Disability-Adjusted Life Years (DALYs) lost. Cancer incidence rose by 33% between 2005 and 2015[28]. Acrylamide is believed to cause cancer by converting to the genotoxic compound glycidamide through the enzyme cytochrome P4502E1[29,30]. While animal studies have shown a dose-response relationship between acrylamide and cancer, particularly in hormone-sensitive organs[31], human studies have mostly found no clear link between dietary acrylamide and cancer, except for a potential increased risk of pancreatic cancer in workers exposed to acrylamide[30]. Some studies have suggested that there is a possible link between dietary acrylamide(AA) intake and a cancer risk such as Oesophageal Cancer a higher risk among overweight or obese individuals with higher AA intake[32], risk Lymphatic malignancies, especially in smoking men[33].

4.2 Neurotoxicity

Acrylamide (AA) is proven to cause neurotoxicity in humans through occupational exposure, leading to peripheral neuropathy symptoms such as numbness and tingling in the limbs. In laboratory animals, chronic AA exposure causes skeletal muscle weakness and ataxia. While the exact mechanism of AA neurotoxicity remains unclear, proposed mechanisms include[34,35]. Impairment of nerve growth occurs when AA interacts with kinesin motor proteins in neurons, disrupting fast anterograde transport of nerve growth factor resulting in death. It also inhibits neurotransmitter uptake into synaptic vesicles, potentially through interactions with sulfhydryl groups[36].

4.3 Reproductive and Developmental Effects

Research on laboratory animals indicates that high doses of AA can cause reproductive toxicity, although no such effects have been documented in humans[25]. In rats, the No Observed Adverse Effect Level (NOAEL) for reproductive toxicity is estimated at 2–5 µg/kg/day, which is four times higher than the doses associated with

neurotoxicity and 2000 times greater than typical dietary exposure[25,37]. In rats, administering 0.5–10 mg/kg of AA resulted in growth retardation, reduced sperm reserves in the epididymis, and testicular lesions. In male rats, doses of 20 mg/kg led to dose-dependent reductions in testosterone and prolactin levels. Similarly, studies on female mice showed that oral exposure to AA decreased body and organ weights, reduced the number of corpora lutea, and lowered serum progesterone levels in a dose-dependent manner[38,39].

5. Method of Detection and Quantification

Acrylamide exposure can be detected by human biomonitoring using either blood[40] or urine[41] Samples. Dietary exposure to acrylamide was assessed by integrating data on its concentrations in foods with food consumption patterns. While a probabilistic approach is recognized as the most accurate method for estimating dietary exposure, a deterministic approach is often suitable as an initial screening tool to evaluate potential risks[42].

Quantification of AA in food has been a need of the hour. It is observed that Novel analytical techniques like capillary electrophoresis, immunoenzymatically test and electrochemical biosensors, are used in the present era in place of traditional methods[43].

Instrumental methods, particularly chromatography, are the most reliable for acrylamide quantification. Techniques like HPLC and the more advanced UPLC offer fast, accurate, and reproducible results, though they demand refined sample preparation and optimized protocols[44]. Gas chromatography (GC), often coupled with mass spectrometry (GC-MS), is widely used for acrylamide analysis, requiring derivatization techniques like bromination or silylation to improve precision and detectability[45].

Biosensors and immunochemical techniques are innovative, cost-effective methods for acrylamide quantification in foods. Biosensors use bioreceptors (e.g., enzymes, antibodies, or microbial cells) to detect acrylamide with high sensitivity and selectivity, even in complex matrices. Types include electrochemical biosensors (e.g., amperometric and potentiometric), which offer advantages like simplicity, rapid assays, and miniaturization for online use. Techniques such as cyclic voltammetry and ion-selective electrodes have enabled detection at trace levels[45,46].

A universal procedure for acrylamide analysis in all food types does not exist due to complex sample preparation and the use of expensive reagents. As a result, only a few laboratories conduct these assays, requiring modern equipment and labour-intensive protocols. The challenges of time-consuming preparation have driven the development of faster, simpler, and cost-effective methods[45].

6. Strategies to Mitigate Formation

Mitigating acrylamide formation in food has become a critical area of research due to its classification as a probable human carcinogen. Strategies to reduce acrylamide are primarily focused on modifying ingredients, optimizing processing conditions, and exploring pre-process interventions[47].

6.1 Modifying Ingredients

Ingredient-level strategies include the use of asparaginase to reduce the precursor asparagine and selecting raw materials with naturally low levels of asparagine and reducing sugars. Process-level strategies emphasize controlling cooking temperatures and durations, as higher thermal loads significantly increase acrylamide formation[48].

6.2 Optimizing Processing Conditions

Techniques such as vacuum baking and frying are promising alternatives, as they operate at lower temperatures, thereby reducing thermal stress while maintaining product quality. Surface treatments like blanching potatoes before frying can also minimize acrylamide precursors. Additionally, combined methods, such as partial conventional baking followed by vacuum baking, have been shown to significantly decrease acrylamide levels by 95% while only vacuum baking reduced it up to 985% and preserved sensory[49].

6.3 Exploring Pre-Process Interventions

There are research article investigates how various soaking pre-treatments (cold and hot water, NaCl, and citric acid solutions) influence the reduction of acrylamide content in French fries prepared under domestic conditions. Soaking pre-treatments, including cold water soaking for

15–120 minutes, reduced acrylamide by 42–89% in pan-fried and up to 47% in fryer-cooked French fries. Hot water blanching at 60–80°C for 5–15 minutes provided even higher reductions, up to 97%, depending on the duration and temperature. Additionally, soaking in NaCl solutions lowered acrylamide by up to 61% by reducing pH and altering heat transfer, while citric acid soaking achieved reductions of up to 97% by significantly lowering pH, inhibiting acrylamide formation[50].

7. Strategies to Reduce Acrylamide Formation

Acrylamide forms in foods primarily through the Maillard reaction, where reducing sugars react with asparagine at temperatures above 120°C, especially in carbohydrate-rich foods like potatoes and cereals[1,2]. High asparagine levels in starchy foods further enhance its production, but pre-cooking methods like blanching or soaking can reduce acrylamide formation by up to 97%[3,4]. Lipid degradation during frying also contributes via reactions with amino acids in fat-rich foods[5]. Factors such as pH, moisture, and starch content in the food matrix significantly influence acrylamide levels, highlighting the importance of controlling cooking conditions and food composition to mitigate its formation[6,7].

Dietary acrylamide exposure mainly comes from high-carbohydrate foods cooked at high temperatures, such as French fries, potato chips, cereal-based products, coffee, baked goods, and toasted bread[8,9]. Darker, more roasted products typically contain higher acrylamide levels[10].

Acrylamide, classified as a probable human carcinogen by the IARC (International Agency for Research on Cancer, 2024)[11], has been linked to cancer in animal studies, particularly in hormone-sensitive organs[12]. While human studies show mixed results, high dietary intake may elevate the risk of cancers such as oesophagal cancer in obese individuals and lymphatic malignancies in smoking men[13,14]. Neurotoxic effects, including peripheral neuropathy, have been observed, especially in occupational exposure, with mechanisms involving disruption of neuronal transport and neurotransmitter uptake[15]. Animal studies also show reproductive toxicity at high doses[16], but human studies have not confirmed similar effects at typical dietary levels[17].

Strategies to reduce acrylamide formation include using asparaginase to lower asparagine levels and selecting low precursor ingredients[18]. Process modifications like vacuum baking and frying reduce acrylamide by minimizing thermal stress, with surface treatments such as blanching potatoes further lowering precursor levels[19,20]. Pre-process interventions, like soaking in hot water or citric acid solutions, have shown significant reductions, with hot water blanching achieving up to 97% reduction[21]. These techniques offer scalable solutions for both domestic and industrial applications[22].

8. Conclusion

Acrylamide formation in food, primarily occurring during high-temperature cooking methods such as frying, roasting, and baking, has raised significant health concerns due to its neurotoxic, genotoxic, and carcinogenic properties[23]. The formation of acrylamide is largely attributed to the Maillard reaction, particularly involving the amino acid asparagine and reducing sugars, with factors such as temperature, cooking time, moisture content, and food composition influencing the extent of its production[24,25]. Potato products, cereals, coffee, and baked goods are some of the major dietary sources of acrylamide[26,27].

Although acrylamide is considered a probable human carcinogen by the International Agency for Research on Cancer (IARC)[28], the link between dietary acrylamide consumption and cancer risk remains controversial, with most studies showing no conclusive evidence of a direct correlation[29,30]. However, acrylamide exposure has been associated with neurotoxicity, and there are concerns about its reproductive and developmental effects based on animal studies[31,32].

To mitigate acrylamide formation, various strategies have been proposed, such as modifying ingredients (e.g., reducing asparagine content), optimizing cooking conditions (e.g., vacuum frying or blanching), and exploring pre-process interventions like soaking in water or acidic solutions[33-35]. While analytical methods like chromatography and emerging biosensors are crucial for detecting and quantifying acrylamide, these methods are expensive and labour-intensive, making them less accessible for widespread use[36,37].

In conclusion, reducing acrylamide levels in foods is a complex challenge that requires the combination of ingredient modifications, optimized cooking processes, and innovative detection techniques[38,39]. Public health measures, including regulatory guidelines and consumer

education, are essential to limit exposure while preserving food quality[40,41].

9. References

1. Bounegru AV, Bounegru I. Acrylamide in food products and the role of electrochemical biosensors in its detection: A comprehensive review. Anal Methods. 2024; 16:2824–2839. https://doi.org/10.1039/D4AY00466C
2. Navruz-Varli S, Mortaş M. Strategies to mitigate acrylamide formation in fried foods. Foods. 2024; 13(4):556. https://doi.org/10.3389/fnut.2023.1297069
3. Tareke E, Rydberg P, Karlsson P, Eriksson S, Törnqvist M. Analysis of acrylamide, a carcinogen formed in heated foodstuffs. J Agric Food Chem. 2002; 50(17):4998–5006. https://doi.org/10.1021/jf020302f
4. Friedman M. Chemistry, biochemistry, and safety of acrylamide: A review. J Agric Food Chem. 2003; 51(16):4504–4526. https://doi.org/10.1021/jf030204+
5. Zhao Y, Long T, Shen L, Huang S, Wu K. Synthesis, characterization, and analysis of the acrylamide- and glycidamide-glutathione conjugates. Chem Biol Interact. 2015; 237:38–46. https://doi.org/10.1016/j.cbi.2015.05.002
6. European Food Safety Authority (EFSA). Scientific opinion on acrylamide in food. EFSA J. 2015; 13(6):4104. https://doi.org/10.2903/j.efsa.2015.4104
7. U.S. Food and Drug Administration (FDA). Guidance for industry on acrylamide in food. Available from: https://www.fda.gov/media/97830/download.
8. Chen C, Song Z, Xu X, Yang X, Wei S, Chen F, *et al.* The neurotoxicity of acrylamide in ultra-processed foods: Interventions of polysaccharides through the microbiota-gut-brain axis. Food Funct. 2024. https://doi.org/10.1039/D4FO03002H
9. Filippini T, Virgolino A. Editorial: Dietary acrylamide in human health. Front Nutr. 2024; 11:1446690. https://doi.org/10.3389/fnut.2024.1446690
10. Mihalache OA, Dall'Asta C. The burden of disease due to dietary exposure to acrylamide in Italy: A risk assessment-based approach. Food Chem Toxicol. 2024; 188(6):114699
11. Augustine DA, Bent GA. Acrylamide, a toxic maillard by-product and its inhibition by sulfur-containing compounds: A mini-review. Front Food Sci Technol. 2022; 2. https://doi.org/10.3389/frfst.2022.1072675
12. Pedreschi F, Granby K, Knudsen D. Acrylamide formation in foods and its reduction methods. Compr Rev Food Sci Food Saf. 2016; 15(4):759–778. https://doi.org/10.1111/1541-4337.12247
13. Huang C, Jiang S, Yang Y, Gao S, Lin Z, Gu W, *et al.* Contaminant acrylamide in food: toxicity, detection methods, mitigation measures and risk assessment for humans. Nat Resour Hum Health. 2021; 1(2):98–109. https://doi.org/10.53365/nrfhh/142376
14. Friedman M, Levin CE. Chemistry, biochemistry, and safety of acrylamide in foods. Compr Rev Food Sci Food Saf. 2019; 18(1):194–224. https://doi.org/10.1111/1541-4337.12439
15. Miller RS, Garcia R. Acrylamide formation and health risks: Lipid-based mitigation strategies. Food Chem. 2016; 199:543–549. https://doi.org/10.1016/j.foodchem.2015.12.069
16. Tsuda T, Fukuoka S. Interaction of lipids and amino acids in acrylamide formation during frying. Food Chem. 2021; 333:127563. https://doi.org/10.1016/j.foodchem.2020.127563
17. Ahn JH, Cho EA, Lee SH. Influence of food matrix on acrylamide formation: The role of starch and water content. J Food Sci. 2018; 83(3):625–631. https://doi.org/10.1111/1750-3841.14046
18. Liu Y, Zhang L, Zhan X. Reducing acrylamide formation in fried foods: A review of recent studies. Food Control. 2015; 54:169–175. https://doi.org/10.1016/j.foodcont.2015.02.021
19. Mottram DS, Wedzicha BL, Dodson AT. Acrylamide is formed in the Maillard reaction. Nature. 2002; 419:448–449. https://doi.org/10.1038/419448a
20. Zhang Y, Bai Y. Acrylamide formation in the Maillard reaction. Food Chem. 2018; 247:239–245. https://doi.org/10.1016/j.foodchem.2017.12.062
21. Asadi S, Aalami M, Shoeibi S, Kashaninejad M, Ghorbani M, Delavar M. Effects of different roasting methods on formation of acrylamide in pistachio. Food Sci Nutr. 2020; 8(6):2875–2881. https://doi.org/10.1002/fsn3.1588
22. Govindaraju I, Sana M, Chakraborty I, Rahman MH, Biswas R, Mazumder N. Dietary Acrylamide: A detailed review on formation, detection, mitigation, and its health impacts. Foods. 2024; 13(4):556. https://doi.org/10.3390/foods13040556
23. Laguzzi F, Filippini T, Virgolino A. Editorial: Dietary acrylamide in human health. Front Nutr. 2024; 11. https://doi.org/10.3389/fnut.2024.1446690
24. Barón Cortés WR, Vásquez Mejía SM, Suárez Mahecha H. Consumption study and margin of exposure of acrylamide in food consumed by the Bogotá population in Colombia. J Food Compos Anal. 2021; 100:103934. https://doi.org/10.1016/j.jfca.2021.103934
25. Zhang Y, Zhang G, Zhang Y. Occurrence and analytical methods of acrylamide in heat-treated foods: Review and recent developments. J Chromatogr A. 2005; 1084(1-2):1–12. https://doi.org/10.1016/j.chroma.2005.03.123
26. Doerge DR, Young JF, McDaniel LP, Twaddle NC, Churchwell MI. Toxicokinetics of acrylamide and glycidamide in Fischer 344 rats. Toxicol Appl Pharmacol.

2005; 208(3):199–209. https://doi.org/10.1016/j.taap.2005.03.003
27. Poteser M, *et al.* Time trends of acrylamide exposure in Europe: Combined analysis of published reports and current HBM4EU studies. Toxics. 2022; 10(8). https://doi.org/10.3390/toxics10080481
28. Kumar J, Das S, Teoh SL. Dietary acrylamide and the risks of developing cancer: Facts to ponder. Front Nutr. 2018; 5:14. https://doi.org/10.3389/fnut.2018.00014
29. Dybing E, *et al.* Human exposure and internal dose assessments of acrylamide in food. Food Chem Toxicol. 2005; 43(3):365–410. https://doi.org/10.1016/j.fct.2004.11.004
30. Hogervorst JG, Schouten LJ, Konings EJ, Goldbohm RA, Van Den Brandt PA. A prospective study of dietary acrylamide intake and the risk of endometrial, ovarian, and breast cancer. Cancer Epidemiol Biomarkers Prev. 2007; 16(11):2304–13. https://doi.org/10.1158/1055-9965.EPI-07-0581
31. Johnson KA, Gorzinski SJ, Bodner KM, *et al.* Chronic toxicity and oncogenicity study on acrylamide incorporated in the drinking water of Fischer 344 rats. Toxicol Appl Pharmacol. 1986; 6(2):172–181. https://doi.org/10.1016/0041-008X(86)90171-6
32. Lujan-Barroso L, *et al.* Dietary intake of acrylamide and oesophagal cancer risk in the European prospective investigation into cancer and nutrition cohort. Cancer Causes Control. 2014; 25(5):639–646. https://doi.org/10.1007/s10552-014-0359-5
33. Bongers ML, Hogervorst JGF, Schouten LJ, *et al.* Dietary acrylamide intake and the risk of lymphatic malignancies: The Netherlands cohort study on diet and cancer. PLoS One. 2012; 7(6):e38016. https://doi.org/10.1371/journal.pone.0038016
34. Spencer PS, Schaumburg HH. A review of acrylamide neurotoxicity Part I: Properties, uses, and human exposure. Can J Neurol Sci. 1974; 1(2):143–150. https://doi.org/10.1017/S0317167100019739
35. McCollister DD, Oyen F, Rowe VK. Toxicology of acrylamide. Toxicol Appl Pharmacol. 1964; (2):172–81. https://doi.org/10.1016/0041-008X(64)90023-1
36. Sickles DW, Brady ST, Testino A, Friedman MA, Wrenn RW. Direct effect of the neurotoxicant acrylamide on kinesin-based microtubule motility. J Neurosci Res. 1996; 46(1):7–17. https://doi.org/10.1002/(SICI)1097-4547(19961001)46:1
37. Wang H, *et al.* Reproductive toxicity of acrylamide-treated male rats. Reprod Toxicol. 2010; 29(2):225–230. https://doi.org/10.1016/j.reprotox.2009.11.002
38. Fatehyab Ali S, Hong J-S, Wilson WE, Uphouse LL, Bondy SC. Effect of acrylamide on neurotransmitter metabolism and neuropeptide levels in several brain regions and upon circulating hormones. Neurotox Res. 1983; 21(3):231–238. https://doi.org/10.1007/BF00317980
39. Wei Q, Li J, Li X, Zhang L, Shi F. Reproductive toxicity in acrylamide-treated female mice. Reprod Toxicol. 2014; 46:121–8. https://doi.org/10.1016/j.reprotox.2014.03.007
40. Schettgen T, Müller J, Fromme H, Angerer J. Simultaneous quantification of haemoglobin adducts of ethylene oxide, propylene oxide, acrylonitrile, acrylamide, and glycidamide in human blood by isotope-dilution GC/NCI-MS/MS. J Chromatogr B. 2010; 878(27):2467–2473. https://doi.org/10.1016/j.jchromb.2009.11.043
41. Heudorf U, Hartmann E, Angerer J. Acrylamide in children: Exposure assessment via urinary acrylamide metabolites as biomarkers. Int J Hyg Environ Health. 2009; 212(2):135–141. https://doi.org/10.1016/j.ijheh.2008.04.006
42. Yu WZ, *et al.* Occurrence and dietary exposure to acrylamide from foods consumed within and outside main meals in Singapore. Foods. 2023; 12(16):3022. https://doi.org/10.3390/foods12163022
43. Oracz J, Nebesny E, Zyzelewicz D. New trends in quantification of acrylamide in food products. Talanta. 2011; 85(1):14–25. https://doi.org/10.1016/j.talanta.2011.08.066
44. Arisseto AP, de Figueiredo Toledo MC, Govaert Y, van Loco J, Fraselle S, Degroodt JM. Modified sample preparation for acrylamide determination in cocoa and coffee products. Food Anal Methods. 2008; 1(1):49–55. https://doi.org/10.1007/s12161-007-9001-4
45. Lee MR, Chang LY, Dou J. Determination of acrylamide in food by solid-phase microextraction coupled to gas chromatography-positive chemical ionization tandem mass spectrometry. Anal Chim Acta. 2007; 582(1):19–23. https://doi.org/10.1016/j.aca.2006.08.042
46. Silva NAF, Matos MJ, Karmali A, Rocha MM. An electrochemical biosensor for acrylamide determination: Merits and limitations. Port Electrochim Acta. 2011; 29(5):361–373. https://doi.org/10.4152/pea.201105361
47. Nguyen KH, Nielsen RH, Mohammadifar MA, Granby K. Formation and mitigation of acrylamide in oven-baked vegetable fries. Food Chem. 2022; 386:132764. https://doi.org/10.1016/j.foodchem.2022.132764
48. Akkurt K, Mogol BA, Gökmen V. Mitigation of acrylamide in baked potato chips by vacuum baking and combined conventional and vacuum baking processes. LWT. 2021; 144:111211. https://doi.org/10.1016/j.lwt.2021.111211
49. Rifai L, Saleh FA. A review on acrylamide in food: Occurrence, toxicity, and mitigation strategies. Toxicol Mech Methods. 2020; 30(4):243–258. https://doi.org/10.1177/1091581820902405
50. Negoiță M, Mihai AL, Horneț GA. Influence of water, NaCl, and citric acid soaking pre-treatments on acrylamide content in French fries prepared in domestic conditions. Foods. 2022; 11(9):1204. https://doi.org/10.3390/foods11091204