Review Article
ISSN (Print): 0971-913X
ISSN (Online): 2455-1244
J Endocrinol Reprod. *Vol 28(3), 159-171, September 2024, DOI: 10.18311/jer/2024/45500*

Dynamic Upstream Controllers of Kisspeptin and its Possible Interactions: A Review Elucidating the Pleiotropic Regulators

Nehareeka Dan, Harsh Shah, Hetvi Shah, Prince Upadhyay, Sripriya Bulusu, A.V. Ramachandran and Parth Pandya*

Department of Biomedical and Lifesciences, School of Science, Navrachana University, Vadodara – 391410, Gujarat, India; parthp@nuv.ac.in

Abstract

Kisspeptin, a neuropeptide initially recognized for its metastasis-suppressing properties, has since emerged as a pivotal regulator in reproductive biology. This review explores the multifaceted roles of kisspeptin, particularly its involvement in the neuroendocrine regulation of puberty, reproduction, and metabolic processes. The Kiss1-Kiss1R signaling pathway is highlighted as crucial for initiating puberty and maintaining reproductive function, primarily through its regulation of Gonadotropin-Releasing Hormone (GnRH) secretion. Additionally, the interactions of kisspeptin with other neuropeptides, such as Agouti-Related Peptide (AgRP), Neuropeptide Y (NPY), and Pro-opiomelanocortin (POMC), emphasize its key role in linking energy balance with reproductive health. The review also discusses upstream regulatory factors, including leptin and melatonin, and their influence on kisspeptin signaling pathways. Moreover, the article delves into the epigenetic regulation of the Kiss1 gene, underscoring its importance in reproductive disorders and potential therapeutic applications. The intricate relationship between kisspeptin and metabolic signals underscores its significance in reproductive endocrinology and opens avenues for addressing metabolic and reproductive disorders. In conclusion, the review suggests future research directions by filling the gap in the existing available data and also elucidates the mechanisms underlying kisspeptin's regulatory functions to develop targeted interventions for related health conditions.

Keywords: Hypothalamic-Pituitary-Gonadal Axis, Kisspeptin, Neuropeptides, Puberty

1. Introduction

The peptide kisspeptin has gained significant attention since its emergence in the scientific community, especially with respect to its involvement in neuroendocrine control of puberty, reproduction, and in cancer biology. It came into light in 1996 when it was first isolated from human melanoma cell line and was known as metastin since it was found to be a potential gene for suppressing metastasis[1]. This discovery happened in the Pennsylvanian town of Hershey, home to the renowned 'Hershey chocolate Kiss', therefore, in its honor, the letters 'KI' were prefixed to suppressor sequence (ss)[2]. Following this discovery, an independent study done in rats in 1999 found an orphan receptor GPR54 belonging to the rhodopsin family, having a 45% resemblance to the galanin receptor family, but no binding affinity[3]. It was later found in 2001 that the kisspeptin gene *kiss1* acted as an endogenous ligand and bound with GPR54[4,5]. As a result, GPR54 was assigned as kisspeptin receptor *kiss1r*. Supporting this, in 2001 four different studies identified kisspeptin to be RF-amide (Arg-Phe-NH_2) peptide natural ligand for GPR54 (kiss1r)[4-7]. Two forms of kisspeptin have since been identified - *kiss1* and *kiss2*.

Interestingly, kisspeptin was initially recognized for its potential in cancer research. Treatment with Kiss1-derived polypeptides reduced metastases in mice engrafted with melanoma cells overexpressing Kiss1R[8]. This shift from cancer research to becoming a cornerstone of reproductive biology highlights kisspeptin's multifaceted significance and the profound impact of its discovery on medical science.

*Author for correspondence

Article Received: 24.08.2024 Accepted on: 10.10.2024

The signaling of Kiss1-Kiss1R gathered the attention of reproductive biologists when in 2003, two independent studies reported that mutations in *kiss1r* resulted in absent or delayed pubertal onset and hypothalamic hypogonadism in humans and mice[9,10]. Similar results were observed in non-mammalian vertebrates like fishes where exogenous injection of kisspeptin externally lead to the initiation of puberty[11,12]. Administration of kisspeptin also resulted in elevated expression of Gonadotropin-Releasing Hormone (GnRH), gonadotropins, and sex hormones[13]. Research on rodents and primates also demonstrated that Kiss1 stimulates the release of GnRH through Kiss1Rs that are expressed in GnRH neurons, furthermore controlling the follicular development and gamete maturation[14,15]. Thus, the notion that kiss1-kiss1r signaling is essential for puberty and reproduction began to take shape.

Puberty is a series of developmental changes that every individual goes through which results in the attainment of reproductive potential and sexual distinction[16]. Several factors have been identified as a modulator of pubertal onset including endogenous signals, environmental cues, and genetic factors, whose dynamic interaction results in the timing of onset of puberty[17,18]. From a neuroendocrine standpoint, puberty starts when there is a complete activation of the Hypothalamic Pituitary Gonadal (HPG) axis[16,18]. It is now a known fact that kisspeptin activates the GnRH to release[19,20], which then stimulates the secretion of gonadotropins Luteinizing Hormone (LH), and Follicle-Stimulating Hormone (FSH). These gonadotropins now act on the gonads which release the sex steroids (estrogen and androgen), reaching full maturity and function during the time of puberty[21].

It is still unclear how kisspeptin gets and decodes the signals from various factors to stimulate the release of GnRH. Therefore, to understand the downstream role of kisspeptin in regulating puberty, reproduction, and sexual maturation, it is important to know its upstream signaling as well. Therefore, the present review focuses on comprehending the various molecules that are involved in activating the kisspeptin signaling pathway.

2. Agouti-related Peptide (AgRP), Pro-opiomelanocortin (POMC) and Neuropeptide Y (NPY)

Agouti-Related Peptide (AgRP) has a direct influence on kisspeptin neurons in the hypothalamus's arcuate nucleus (ARC), linking energy balance with reproductive function, as noted by Merkley *et al.*[22]. This study revealed that many kisspeptin neurons express Melanocortin 3 and 4 receptors (MC3R and MC4R), which are involved in AgRP's actions, suggesting an inhibitory effect of AgRP on reproductive activities. While the anatomical basis for AgRP's interaction with kisspeptin is established, the exact functional outcomes of this interaction, particularly how it affects hormone secretion like LH, require further exploration.

Pro-opiomelanocortin (POMC) neurons also interact with kisspeptin neurons, suggesting a regulatory role in reproductive and metabolic processes, though the specifics of this interaction remain less defined[23]. It is not entirely clear whether the influence of POMC on kisspeptin is direct or indirect, or how significant this interaction is for overall metabolic and reproductive health.

Neuropeptide Y (NPY), another key player in the reproductive axis, has been shown to influence kisspeptin activity. Studies like those by Belsham *et al.*[24] and Backholer *et al.*,[25] demonstrate that kisspeptin directly regulates NPY synthesis and secretion in hypothalamic cell lines, potentially impacting GnRH secretion. This suggests that NPY might play a role in the broader regulatory network of reproductive function, although its exact role in conditions like Polycystic Ovary Syndrome (PCOS) remains under investigation.

Recent research also highlights the effects of obesity on kisspeptin function, particularly in terms of reproductive hormone regulation. Azizi *et al.*,[26] found that obesity leads to a suppression of kisspeptin neurons, leading to lower levels and reduced pulse frequency of LH in obese mice despite normal pituitary responsiveness. This suppression is linked to reduced glutamate receptor expression and diminished interaction with anorexigenic POMC neurons, elucidating potential mechanisms by which obesity impairs GnRH secretion.

Adding to the understanding of neuropeptide interactions with kisspeptin, Villa *et al.*[27] explored the role of NPY in the expression of Kiss1 and Gpr54 genes in the pituitary gland. They discovered that NPY stimulates the expression of these genes, an effect that can be blocked by the receptor antagonist BIBP3226, highlighting the complex regulatory network involving neuropeptides in reproductive hormone control. Collectively, these studies emphasize the intricate and significant influence of neuropeptides like AgRP, POMC, and NPY on kisspeptin regulation and, consequently, on reproductive

health. Understanding these interactions is crucial for comprehending how metabolic states affect reproductive functions and for developing targeted interventions for reproductive and metabolic disorders. Further research is necessary to clarify the precise mechanisms and functional consequences of these neuropeptide interactions on reproductive health.

3. Irisin

The discovery of novel hormones and peptides has revolutionized our understanding of metabolic and reproductive physiology. Among these, irisin and kisspeptin have garnered significant attention due to their multifaceted roles in energy homeostasis, reproductive function, and overall metabolic regulation. Irisin, a myokine released during physical exercise, and kisspeptin, a neuropeptide critical for the initiation of puberty and regulation of fertility, have shown intriguing interactions that offer potential therapeutic implications for metabolic and reproductive disorders. Irisin was identified as a muscle-secreted hormone that converts white adipose tissue into brown-like fat, enhancing thermogenesis and energy expenditure. This myokine is cleaved from the fibronectin Type III domain-containing protein 5 (FNDC5) and has been shown to play a pivotal role in metabolic processes, including glucose homeostasis and lipid metabolism. Studies have demonstrated that irisin levels increase with physical activity, highlighting its role in mediating the beneficial effects of exercise on metabolic health[28]. Additionally, irisin's influence on the central nervous system indicates that it may affect brain function and neuroprotection, though these areas require further exploration.

On the other hand, kisspeptin's involvement in the modulation of metabolic processes has been suggested, linking reproductive health to metabolic status. The hormone's potential therapeutic applications include treating disorders such as hypogonadotropic hypogonadism and PCOS, where kisspeptin administration has shown promising results in restoring reproductive function[29]. The interaction between irisin and kisspeptin presents a fascinating area of research, particularly concerning their combined effects on metabolic and reproductive health. Studies suggest that kisspeptin may influence irisin release, thereby linking reproductive signals to metabolic responses[30]. For instance, a study found that peripheral administration of kisspeptin increased circulating levels of irisin in humans, indicating a possible therapeutic avenue for metabolic disorders through the modulation of these hormones[30].

The role of irisin in the hypothalamus, where it may interact with kisspeptin neurons, highlights its potential impact on the HPG axis and reproductive function. This interaction suggests that irisin could modulate GnRH release, thereby influencing LH and FSH secretion, integrating energy balance with reproductive health[31]. Understanding the mechanisms underlying the interaction between irisin and kisspeptin could lead to integrated therapeutic strategies that address both metabolic and reproductive health.

4. Neurokinin-B (NKB) and Dynorphin (DYN) Network

Kisspeptin neurons influence GnRH secretion with permissive or opposing signals mediated by NKB and DYN, forming the Kisspeptin-NKB-DYN (KNDy) neuronal system. Some of the recent investigations have described the involvement of Kiss1, NKB, and DYN in KNDy neurons for regulating pulsatile GnRH secretion. Substantial evidence supports the role of these peptides in generating GnRH pulses in female mammals, though direct proof has been challenging due to difficulties with gene-edited animal models[32-34]. Recently, Kiss1 knockout rats have established the physiological functions of kisspeptin, for both pulsatile and surge-mode GnRH release, because these rats exhibited hypogonadal phenotypes and almost absent pulsatile LH secretion[35-38].

To address this issue, one study demonstrates the restoration of Kiss1 expression, exclusively in the KNDy neurons of global Kiss1 knockout rats. This rescue partially revived pulsatile LH discharge, with the extent of restoration proportional to the number of KNDy neurons preserved. While resuscitating KNDy neurons culminated in follicular development, it failed to offer optimal recovery of ovulation. This elicited that despite its importance in GnRH/LH pulse generation, KNDy neurons do not solely dictate surge generation[39].

Further tests on conditional ARC Kiss1 knockdown rats showed that ARC Kiss1 neurons are crucial for generating GnRH pulses, as nearly 90% of Kiss1-expressing cells were knocked down, leading to a complete loss of pulsatile LH secretion. This highlights the essential role of ARC Kiss1 neurons in regulating reproductive hormone secretion. Based on these results the authors state that there must be functional overlap among the

KNDy neurons, with a specific subpopulation capable of regulating GnRH pulses and folliculogenesis.

Moreover, although NKB and DYN overlap with GnRH fibers, they possibly do not regulate the pulsatile release of GnRH by instantly influencing the neurons[40,41]. As discussed until now, the possible interplay between these neuropeptides and the hormones in triggering the initiation of the HPG axis has been shown in Figure 1.

5. Melatonin: The Photo-regulator of Kisspeptin

Melatonin, a neuroendocrine hormone, is primarily secreted by the pineal gland. This hormone follows a circadian rhythm regulated by the suprachiasmatic nucleus, with secretion levels peaking at night in response to changes in light and dark periods in nature and regulates the reproductive axis across different species including fish, rats, and humans, through the modulation of kisspeptin[42-44]. Reiter *et al*., demonstrated that melatonin acts through G protein-coupled receptors, chiefly M1 and M2, distributed in the central nervous system[45]. Another study found that melatonin regulates the key enzyme involved in kisspeptin biosynthesis by modulating transcription factors such as CREB (cAMP response element-binding protein)[44]. Additionally, melatonin's regulatory effects on kisspeptin are influenced by photoperiod, as melatonin secretion is tightly controlled by the light-dark cycle, affecting seasonal breeding patterns in fish (Figure 2)[46]. In teleost, Kiss neurons are directly regulated by pituitary endocrine cells, rather than acting through GnRH neurons, suggesting melatonin involvement in the control of kiss expression[47,48]. A study by Carnevali *et al*., demonstrated that melatonin exposure via immersion increased mRNA transcript levels of *kiss1* and *kiss2* in the brain and *lhb* in the pituitary of adult female zebrafish[49]. In contrast, in male Sea bass, prolonged exposure to melatonin *via* implants caused decreased brain mRNA levels of *kiss1* and *kiss2*[50].

In contrast to fish, short photoperiod upregulates both ARC Kiss1 mRNA and protein in winter breeding sheep, increasing the amount of ARC Kiss neurons and




Figure 1. The image showcases the possible interplay of neuropeptides (NKB, DYN, AgRP, NPY, POMC, Kiss1, GnRH) and hormones (leptin, LH, FSH) that leads to the activation of the HPG axis.
The solid lines represent activation, dashed arrows show indirect activation and the blurred end lines show the inhibitory role. During the transition of pubertal onset, the leptin levels are found to be elevated, which in turn leads to inhibition of AgRP and NPY. However, leptin can possibly be a stimulator of POMC from POMC neurons. NKB and DYN are coexpressed with kisspeptin, forming the KNDy neurons, which further stimulates the Gnrh neurons to release GnRH in pulsatile form. Gnrh in turn activates the HPG axis to further downstream signaling and release of LH and FSH.

synaptic connections across Kiss and GnRH neurons[51]. Whereas in summer breeders, melatonin inhibits the activity of KISS neurons[44]. For example, a study by Revel *et al.*, and Ansel *et al.*, demonstrated that using a combination of photoperiod manipulation, pinealectomy, and melatonin administration, melatonin significantly downregulated ARC Kiss1 mRNA in Syrian hamsters, and the effect was further modulated by negative steroid feedback[50-52]. Likewise, Turkish Hamsters (*Mesocricetus brandti*) and Striped Hamsters (*Cricetulus barabensis*) showed inhibitory effects of melatonin on Kiss1 mRNA. Due to the long photoperiod, ARC Kiss1 mRNA levels were reduced, as substantial negative sex steroid feedback overcame the melatonin signal[53,54]. Castration in the long photoperiod restored the levels of both Kiss1 and Kiss2[55]. A study by Li *et al.*, suggests that melatonin does not directly regulate kisspeptin in ovine ARC[56]. Instead, melatonin influences Kisspeptin expression indirectly through its effects on other regulatory factors, such as prolactin[57]. Putting the existing studies together, it can be said that melatonin has a direct or indirect role in stimulating the activity of kisspeptin[58]. Further research is needed to clarify the interactive roles of melatonin and kisspeptin and their impact on pubertal onset.

6. Leptin

White adipose tissue is the primary energy storage site, with adipocytes secreting the anorexigenic adipokine leptin[59]. Circulating leptin levels reflect body fat mass and act as a metabolic signal to the central neuroendocrine regulatory centers of appetite, energy balance, and metabolism[60]. As Kiss1 neurons have a significant role in the neuroendocrine control of reproduction, they are plausible conveyors of metabolic impact on GnRH neurons. However, data suggests that Kiss1 neurons may be involved in this integrative process without directly targeting these metabolic signals, as they lack leptin receptors[61,62]. Various studies also exhibited that in mice and rhesus macaques, after 48 hours of fasting, downregulation in the hypothalamic transcript levels of both Kiss1 and Kiss1r were noted, in contrast to rats[63-65]. Metabolic hormones such as leptin and insulin directly trigger POMC neurons, which in turn excites Kisspeptin




Figure 2. The image represents a possible interaction between melatonin and kisspeptin in regulating the HPG axis. Before the onset of puberty, the melatonin levels are high which inhibits the gonadostat from getting activated. But during the onset of puberty, the melatonin level decreases which might be responsible for kisspeptin to activate the gonadostat and stimulates the release of GnRH, and thereby initiating the axis.

neurons, signifying an important and crucial molecular pathway linking metabolism, food intake, and kisspeptin regulation[66-68].

7. Steroidal Control on Kisspeptin in ARC and AVPV

In the rodent brain, kisspeptin neurons are concentrated into two distinct neuronal populations in the hypothalamus, namely, ARC and anteroventral periventricular nucleus (AVPV)[69,70]. Interestingly, both populations, despite being involved in metabolism and reproduction, show a contrast in Kiss1 mRNA expression[61,71-73]. Prior studies have investigated kisspeptin-dependent GnRH release, which subsequently triggers the release of pituitary hormones LH and FSH, functioning within the HPG axis[74]. Interestingly, GnRH neurons lack sex steroid receptors such as estrogen, whereas, kisspeptin neurons in the ARC and AVPV have estrogen receptor-α (ERα)[75,76]. Additionally, kisspeptin neurons project their axons to the GnRH neurons in the hypothalamus[77]. Therefore, an indirect link between sex steroids and GnRH release is established *via* the stimulation of kisspeptin neurons in the hypothalamus. In the AVPV, estradiol positively regulates kiss1 expression, as shown in Figure 3 [78]. During the preovulatory phase, estradiol levels rise and bind to ERα in the AVPV, causing the depolarization of these neurons and subsequent release of kisspeptin[79]. This cascades down the process to the hypothalamus and GnRH is released into circulation. Here, an unceasing stimulation by estrogen leads to GnRH neurons being constantly depolarised and thus the pituitary hormone, LH levels drastically increase. Consequently, this increase leads to an estrogen-dependent LH surge in female rodents.

On the other hand, the ARC region in the rodent hypothalamus houses the KNDy neurons[80]. Estrogen inhibits Kiss1 expression in pre and post-pubertal female rats[81]. The 'gonadostat hypothesis' supports this regulatory pathway, suggesting that during pubertal transition, sex steroids are in control of the hypothalamic and pituitary secretions[82]. Reduced sex steroid levels during puberty diminish this feedback inhibition, leading to increased GnRH and gonadotropin levels[82]. Moreover, the 'KNDy Hypothesis' was postulated to summarise GnRH or Gonadotropin pulse generation[83]. KNDy neurons synchronize their activity to produce pulsatile GnRH release[84]. Neurokinin B initiates the trigger to release kisspeptin, leading to a sharp increase in GnRH and with it, LH levels. Next, the Dynorphin signal stops the pulse and lastly, kisspeptin is considered as the final output signal[80]. Surprisingly, certain studies have demonstrated the pharmacologic intervention of genes encoding for NKB, DYN, and their receptors, NK3R and K-Opioid Receptor (KOR), respectively, does not necessarily diminish the GnRH pulse generation altogether[85]. Therefore, this hypothesis is under further investigation to establish the distinct role of KNDy neurons.

Kisspeptin expression shows sexual dimorphism in mammals. Paraventricular nucleus (PV) is highly prominent in females compared to males; therefore, kisspeptin regulation by estrogen is highly evident in females[86]. In contrast, it was demonstrated that in males, the sex steroid testosterone levels were highly affected due to KissR knockout, resulting in female-like characters in male rodents[86,87]. This highlights the relevance of the Kisspeptin-GnRH circuit during sexual development and differentiation.

To summarise, Kiss1 mRNA expression differs in mammalian hypothalamic neuronal populations under the influence of sex steroids. Kisspeptin neurons of AVPV and ARC regions of the hypothalamus express Erα that, upon estradiol stimulation, either results in histone acetylation or deacetylation of the Kiss1 promoter, respectively. This change results in the distinct pulsatile and surge release activity of GnRH or gonadotropins that regulate reproductive roles in mammals.

8. Epigenetic Regulators of Kiss1

Epigenetic modifications play a crucial role in regulating Kiss1 gene expression in the context of reproductive disorders. Studies have shown that epigenetic mechanisms, such as DNA methylation, histone modifications, and chromatin remodeling, are involved in the regulation of Kiss1 expression as shown in Figure 4[88-90]. These modifications can impact the function of kisspeptin neurons in the hypothalamus, affecting reproductive processes like puberty onset, maintenance of adult reproductive function, and the development of neuroendocrine disorders associated with aging[88,91]. Furthermore, epigenetic changes in the Kiss1 have been linked to metabolic disorders like Type 2 diabetes mellitus, highlighting the intricate interplay between epigenetic regulation and reproductive health in the context of metabolic diseases[89]. Understanding and targeting these




Figure 3. Effect of estradiol or E2 on kisspeptin release in hypothalamic neuronal populations, namely, AVPV and ARC. E2 binds to Estrogen receptor-α (Erα), present on the Kiss1 neuron in AVPV, and induces the transcription of Kiss1 mRNA. Upon translation, kisspeptin is released into the synapse and binds to its receptor GPR54 on GnRH neurons in the Hypothalamus. Kisspeptin depolarises the neuron and thus induces GnRH into circulation. Upon reaching the anterior pituitary, GnRH binds to its receptor GnRH-R, stimulating the release of gonadotropins, LH, and FSH. Finally, estradiol synthesis is promoted in the ovary, which sends feedback to the hypothalamus. Due to the positive feedback, the AVPV region observes an LH surge as shown in Figure 3. Conversely, in the ARC, ERα found on KNDy neurons inhibits the Kiss1 mRNA by deacetylation of its promoter. Therefore, a pulsatile release of GnRH or gonadotropins is observed.

epigenetic modifications could offer new therapeutic avenues for managing reproductive disorders associated with kisspeptin dysregulation. In female reproductive aging, hypoacetylation of histone H3 in the AVPV leads to decreased Kiss1 expression and LH surge dysfunction. Studies have explored the epigenetic regulation of Kiss1 and Kiss1r genes in the hypothalamus at puberty onset, revealing that methylation of these genes can alter reproductive processes during puberty[92,93]. The epigenetic transition of the Kiss1 gene from transcriptional repression to activation is associated with the Polycomb Group (PcG) repression mechanisms involving Mixed-Lineage Leukaemia 1 (MLL1) and 3 (MLL3), members of the Trithorax Group (TrxG) of chromatin modifiers. MLL1 modifies the chromatin structure at Kiss1 and Tac3 promoters from a repressive to a permissive state, while MLL3 activates a Kiss1 enhancer from a poised to an active state[94].

The activation of the Kiss1 gene in the ARC at puberty is the result of EED protein loss from the Kiss1 promoter and activating histone H3 changes such as H3K4 trimethylation and H3K9/14 acetylation[94]. SIRT1, which is highly expressed in Kiss1 neurons in the ARC, interacts with the PcG complex to repress the Kiss1 promoter through repressive histone modifications. During Puberty, SIRT1 is removed from the Kiss1 promoter, leading to the

activation of Kiss1 transcription and the onset of puberty. Undernutrition delays puberty by maintaining SIRT1 at the Kiss1 promoter, while overnutrition accelerates puberty by promoting the premature removal of SIRT1 from the promoter.

9. Future Scope

Despite the significant progress in understanding kisspeptin's functions, many questions remain unanswered, particularly concerning the upstream regulatory mechanisms and the epigenetic modulation of the Kiss1 gene. Emerging research has also highlighted the role of microRNAs (miRNAs) in the regulation of kisspeptin expression, adding another layer of complexity to its regulatory network[95]. miRNAs, which are short non-coding RNAs, have been shown to modulate gene expression post-transcriptionally and are now recognized as crucial regulators of kisspeptin signaling. The miRNA-mediated regulation of kisspeptin could provide new insights into how its expression is finely tuned in response to various physiological and environmental cues. Understanding this miRNA-mediated regulation could also open up new therapeutic avenues, particularly in the context of reproductive and metabolic disorders where kisspeptin dysregulation plays a key role. Additionally, the potential of kisspeptin as a therapeutic agent in treating infertility and other reproductive conditions deserves further exploration, particularly in clinical settings.

Another promising avenue for future research is the exploration of kisspeptin's role in sexual differentiation and the development of secondary sexual characteristics[96]. Investigating how kisspeptin signaling varies across different stages of life and in different physiological conditions could provide deeper insights into its broader biological functions. Furthermore, the development of novel kisspeptin analogs or antagonists could offer new therapeutic options for managing conditions related to puberty and reproductive health.




Figure 4. Demonstrates how epigenetic mechanisms, including DNA methylation, histone modifications, and chromatin remodeling, regulate the expression of the Kiss1 gene.
The image shows the epigenetic regulation of the Kiss1 gene is a multi-layered process involving DNA methylation, histone modifications, and chromatin remodeling. These processes work together to control the expression of the Kiss1 gene, which is crucial for reproductive function. DNA methylation at the Kiss1 promoter, facilitated by DNMTs, typically represses gene expression. In parallel, histone modifications such as acetylation and methylation are orchestrated by enzymes including HATs, HDACs, and HDMs, which alter the chromatin structure. Proteins like MLL1 and MLL3 play a pivotal role in chromatin remodeling by converting repressive chromatin into a more open and permissive state, enabling the activation of Kiss1 during key stages such as puberty.

10. Conclusion

The research on kisspeptin has significantly advanced our understanding of reproductive biology and neuroendocrine regulation, shedding light on its critical role in the onset of puberty, reproductive function, and metabolic processes. Kisspeptin's discovery as a key regulator of GnRH secretion has positioned it as a cornerstone of reproductive endocrinology, offering insights into the complex interplay between the central nervous system and reproductive organs. The peptide's interactions with other neuropeptides, such as AgRP, NPY, and POMC, further demonstrate its central role in integrating metabolic and reproductive functions, particularly under conditions of energy balance or imbalance.

The exploration of these areas holds the potential to uncover new therapeutic targets for reproductive disorders, such as hypogonadotropic hypogonadism, PCOS, and other conditions associated with kisspeptin dysregulation. Moreover, the role of kisspeptin in metabolic regulation suggests its broader implications in addressing metabolic disorders like obesity and Type 2 diabetes, where energy homeostasis is closely linked to reproductive health.

In nutshell, while significant strides have been made in understanding kisspeptin's role in reproductive biology, much remains to be explored. The continued investigation into its regulatory mechanisms, including miRNA-mediated regulation, therapeutic potential, and broader biological functions will likely yield important discoveries, contributing to both basic science and clinical medicine. Kisspeptin's multifaceted role in connecting metabolic and reproductive health underscores its importance as a key target for future research and therapeutic development.

11. References

1. Lee JH, Miele ME, Hicks DJ, *et al*. KiSS-1, a novel human malignant melanoma metastasis-suppressor gene. J Natl Cancer Inst. 1996; 88(23):1731-1737. https://doi.org/10.1093/jnci/88.23.1731
2. Oakley AE, Clifton DK, Steiner RA. Kisspeptin signaling in the brain. Endocr Rev. 2009; 30(6):713-743. https://doi.org/10.1210/er.2009-0005
3. Lee DK, Nguyen T, O'Neill GP, *et al*. Discovery of a receptor related to the galanin receptors. FEBS Lett. 1999; 446(1):103-107. https://doi.org/10.1016/S0014-5793(99)00009-5
4. Ohtaki T, Shintani Y, Honda S, *et al*. Metastasis suppressor gene KiSS-1 encodes peptide ligand of a G-protein-coupled receptor. Nature. 2001; 411(6837):613-617. https://doi.org/10.1038/35079135
5. Kotani M, Detheux M, Vandenbogaerde A, *et al*. The metastasis suppressor gene KiSS-1 encodes Kisspeptins, the natural ligands of the Orphan G protein-coupled receptor GPR54. J Biol Chem. 2001; 276(37):34631-34636. https://doi.org/10.1074/jbc.M104847200
6. Muir AI, Chamberlain L, Elshourbagy NA, *et al*. AXOR12, a novel human G protein-coupled receptor, activated by the peptide KiSS-1. J Biol Chem. 2001; 276(31):28969-28975. https://doi.org/10.1074/jbc.M102743200
7. Clements MK, McDonald TP, Wang R, *et al*. FMRF amide-related neuropeptides are agonists of the Orphan G-protein-coupled receptor GPR54. Biochem Biophys Res Commun. 2001; 284(5):1189-1193. https://doi.org/10.1006/bbrc.2001.5098
8. Navenot JM, Evans B, Oishi S, *et al*. The metastasis suppressor KISS1 lacks antimetastatic activity in the C8161.9 xenograft model of melanoma. Melanoma Res. 2012; 22(2):140-150. https://doi.org/10.1097/CMR.0b013e328350fa07
9. de Roux N, Genin E, Carel JC, *et al*. Hypogonadotropic hypogonadism due to loss of function of the KiSS1-derived peptide receptor GPR54. Proc Natl Acad Sci. 2003; 100(19):10972-10976. https://doi.org/10.1073/pnas.1834399100
10. Seminara SB, Messager S, Chatzidaki EE, *et al*. The GPR54 gene as a regulator of puberty. N Engl J Med. 2003; 349(17):1614-1627. https://doi.org/10.1056/NEJMoa035322
11. Rather MA, Bhat IA, Gireesh-Babu P, *et al*. Molecular characterisation of kisspeptin gene and effect of nano-encapsulted kisspeptin-10 on reproductive maturation in *Catla catla*. Domest Anim Endocrinol. 2016; 56:36-47. https://doi.org/10.1016/j.domaniend.2016.01.005
12. Fatsini E, Fernandez I, Anjos C, *et al*. Kisspeptin influences the reproductive axis and circulating levels of microRNAs in senegalese sole. Int J Mol Sci. 2020; 21(23):9051. https://doi.org/10.3390/ijms21239051
13. Park JW, Jin YH, Oh SY, *et al*. Kisspeptin2 stimulates the HPG axis in immature *Nile tilapia* (*Oreochromis niloticus*). Comp Biochem Physiol B Biochem Mol Biol. 2016; 202:31-38. https://doi.org/10.1016/j.cbpb.2016.07.009
14. Shahab M, Mastronardi C, Seminara SB, *et al*. Increased hypothalamic GPR54 signaling: A potential mechanism for initiation of puberty in primates. Proc Natl Acad Sci. 2005; 102(6):2129-2134. https://doi.org/10.1073/pnas.0409822102
15. Irwig MS, Fraley GS, Smith JT, *et al*. Kisspeptin activation of gonadotropin releasing hormone neurons and regulation

of KiSS-1 mRNA in the male rat. Neuroendocrinol. 2005; 80(4):264-272. https://doi.org/10.1159/000083140
16. Grumbach MM. The neuroendocrinology of human puberty revisited. Horm Res. 2002; 57(2):2-14. https://doi.org/10.1159/000058094
17. Parent AS, Teilmann G, Juul A, *et al*. The timing of normal puberty and the age limits of sexual precocity: Variations around the world, secular trends, and changes after migration. Endocr Rev. 2003; 24(5):668-693. https://doi.org/10.1210/er.2002-0019
18. Ojeda S, Skinner M. Puberty in the rat. Phys Reprod. 2006; 2:2061-2126. https://doi.org/10.1016/b978-012515400-0/50043-9
19. Tassigny XA, Fagg LA, Carlton MBL, *et al*. Kisspeptin can stimulate Gonadotropin-Releasing Hormone (GnRH) release by a direct action at GnRH nerve terminals. Endocrinol. 2008; 149(8):3926-3932. https://doi.org/10.1210/en.2007-1487
20. Okamura H, Tsukamura H, Ohkura S, *et al*. Kisspeptin and GnRH pulse generation. Adv Exp Med Biol. 2013; 784:297-323. https://doi.org/10.1007/978-1-4614-6199-9_14
21. Spaziani M, Tarantino C, Tahani N, *et al*. Hypothalamo-pituitary axis and puberty. Mol Cell Endocrinol. 2021; 520:111094. https://doi.org/10.1016/j.mce.2020.111094
22. Merkley CM, Shuping SL, Sommer JR, *et al*. Evidence That agouti-related peptide may directly regulate Kisspeptin neurons in male sheep. Metabolites. 2021; 11(3):138. https://doi.org/10.3390/metabo11030138
23. Ann J, Li Q, Millar RP, *et al*. Kisspeptin signaling is required for the luteinizing hormone response in anestrous ewes following the introduction of males. Plos One. 2013; 8(2):e57972-e57972. https://doi.org/10.1371/journal.pone.0057972
24. Kim GL, Dhillon SS, Belsham DD. Kisspeptin directly regulates neuropeptide Y synthesis and secretion *via* the ERK1/2 and p38 mitogen-activated protein kinase signaling pathways in NPY-secreting hypothalamic neurons. Endocrinology. 2010; 151(10):5038-5047. https://doi.org/10.1210/en.2010-0521
25. Backholer K, Smith JT, Rao A, *et al*. Kisspeptin cells in the ewe brain respond to leptin and communicate with neuropeptide Y and proopiomelanocortin cells. Endocrinology. 2010; 151(5):2233-2243. https://doi.org/10.1210/en.2009-1190
26. Azizi V, Oryan S, Khazali H, *et al*. Upregulation of Kiss-1 and Gpr54 genes expression in pituitary of male rats following the central administration of neuropeptide Y. Int J Basic sci Med. 2019; 4(4):137-142. https://doi.org/10.34172/ijbsm.2019.03
27. Villa PA, Ruff RER, Jamieson BB, *et al*. Obesity alters POMC and Kisspeptin neuron cross talk leading to reduced luteinizing hormone in male mice. J Neurosci. 2024; 44(28):e0222242024. https://doi.org/10.1523/jneurosci.0222-24.2024
28. Bostrom P, Wu J, Jedrychowski MP, *et al*. A PGC1-α-dependent myokine that drives brown-fat-like development of white fat and thermogenesis. Nature. 2012; 481(7382):463-468. https://doi.org/10.1038/nature10777
29. Chan YM, Fingert SB, Wong KM, *et al*. Kisspeptin/Gpr54-independent gonadotrophin-releasing hormone activity in Kiss1 and Gpr54 mutant mice. J Neuroendocrinol. 2009; 21(12):1015-1023. https://doi.org/10.1111/j.1365-2826.2009.01926.x
30. Shamas S. Changes in Irisin release in response to peripheral Kisspeptin-10 administration in healthy and obese adult men. Acta Endo (Buc). 2019; 15(3):283-288. https://doi.org/10.4183/aeb.2019.283
31. Sengupta P, Dutta S, Karkada IR, *et al*. Irisin, energy homeostasis and male reproduction. Front Physiol. 2021; 12. https://doi.org/10.3389/fphys.2021.746049
32. Uenoyama Y, Nakamura S, Hayakawa Y, *et al*. Lack of pulse and surge modes and glutamatergic stimulation of luteinising hormone release in Kiss1 knockout rats. J Neuroendocrinol. 2015; 27(3):187-197. https://doi.org/10.1111/jne.12257
33. Clarkson J, Han SY, Piet R, *et al*. Definition of the hypothalamic GnRH pulse generator in mice. PNAS. 2017; 114(47):E10216-E10223. https://doi.org/10.1073/pnas.1713897114
34. Han SY, McLennan T, Czieselsky K, *et al*. Selective optogenetic activation of arcuate kisspeptin neurons generates pulsatile luteinising hormone secretion. PNAS. 2015; 112(42):13109-13114. https://doi.org/10.1073/pnas.1512243112
35. Wakabayashi Y, Nakada T, Murata K, *et al*. Neurokinin B and Dynorphin A in Kisspeptin neurons of the Arcuate nucleus participate in generation of periodic oscillation of neural activity driving pulsatile gonadotropin-releasing hormone secretion in the goat. J Neurosci. 2010; 30(8):3124-3132. https://doi.org/10.1523/JNEUROSCI.5848-09.2010
36. Ohkura S, Takase K, Matsuyama S, *et al*. Gonadotrophin-releasing hormone pulse generator activity in the hypothalamus of the goat. J Neuroendocrinol. 2009; 21(10):813-821. https://doi.org/10.1111/j.1365-2826.2009.01909.x
37. Goodman RL, Hileman SM, Nestor CC, *et al*. Kisspeptin, Neurokinin B, and Dynorphin act in the arcuate nucleus to control activity of the GnRH pulse generator in Ewes. Endocrinology. 2013; 154(11):4259-4269. https://doi.org/10.1210/en.2013-1331
38. Lehman MN, He W, Coolen LM, *et al*. Does the KNDy model for the control of gonadotropin-releasing hormone pulses apply to monkeys and humans? SeminReprod Med. 2019; 37(02):71-83. https://doi.org/10.1055/s-0039-3400254
39. Nagae M, Uenoyama Y, Okamoto S, *et al*. Direct evidence that KNDy neurons maintain gonadotropin pulses and folliculogenesis as the GnRH pulse generator. Proc

Natl Acad Sci. 2021; 118(5):e2009156118. https://doi.org/10.1073/pnas.2009156118
40. Popa SM, Moriyama RM, Caligioni CS, *et al*. Redundancy in Kiss1 expression safeguards reproduction in the mouse. Endocrinol. 2013; 154(8):2784-2794. https://doi.org/10.1210/en.2013-1222
41. Krajewski SJ, Anderson MJ, Iles-Shih L, *et al*. Morphologic evidence that neurokinin B modulates gonadotropin-releasing hormone secretion *via* neurokinin 3 receptors in the rat median eminence. J Comp Neurol. 2005; 489(3):372-386. https://doi.org/10.1002/cne.20626
42. Shao R, Wang Y, He C, *et al*. Melatonin and its emerging physiological role in reproduction: A review and update. Curr Mol Med. 2024; 24(4):449-456. https://doi.org/10.2174/1566524023666230417103201
43. Menendez JM, Campa CM. Melatonin: An anti-tumor agent in hormone-dependent cancers. Int J Endocrinol. 2018; 2018:1-20. https://doi.org/10.1155/2018/3271948
44. Ciani E, Haug TM, Maugars G, *et al*. Effects of melatonin on anterior pituitary plasticity: A comparison between Mammals and Teleosts. Front Endocrinol (Lausanne). 2021; 11. https://doi.org/10.3389/fendo.2020.605111
45. Reiter RJ, Tan DX, Galano A. Melatonin: Exceeding expectations. Physiol. 2014; 29(5):325-333. https://doi.org/10.1152/physiol.00011.2014
46. Falcon J, Besseau L, Sauzet S, *et al*. Melatonin effects on the hypothalamo-pituitary axis in fish. Trends Endocrin Met. 2007; 18(2):81-88. https://doi.org/10.1016/j.tem.2007.01.002
47. Dufour S, Querat B, Tostivint H, *et al*. Origin and evolution of the neuroendocrine control of reproduction in vertebrates, with special focus on genome and gene duplications. Physiol Rev. 2020; 100(2):869-943. https://doi.org/10.1152/physrev.00009.2019
48. Pasquier J, Kamech N, Lafont AG, *et al*. Molecular evolution of GPCRS: Kisspeptin or kisspeptin receptors. J Mol Endocrinol. 2014; 52(3):T101-T117. https://doi.org/10.1530/JME-13-0224
49. Carnevali O, Gioacchini G, Maradonna F, *et al*. Melatonin induces follicle maturation in *Danio rerio*. Plos One. 2011; 6(5):e19978-e19978. https://doi.org/10.1371/journal.pone.0019978
50. Revel FG, Ansel L, Klosen P, *et al*. Kisspeptin: A key link to seasonal breeding. Rev Endocr Metab Disord. 2007; 8(1):57-65. https://doi.org/10.1007/s11154-007-9031-7
51. Smith JT, Coolen LM, Kriegsfeld LJ, *et al*. Variation in Kisspeptin and RFamide-Related Peptide (RFRP) expression and terminal connections to gonadotropin-releasing hormone neurons in the brain: A novel medium for seasonal breeding in the sheep. Endocrinol. 2008; 149(11):5770-5782. https://doi.org/10.1210/en.2008-0581
52. Ansel L, Bolborea M, Bentsen AH, *et al*. Differential regulation of Kiss1 expression by melatonin and gonadal hormones in male and female *Syrian Hamsters*. J Biol Rhythms. 2010; 25(2):81-91. https://doi.org/10.1177/0748730410361918
53. Piekarski DJ, Jarjisian SG, Perez L, *et al*. Effects of pinealectomy and short day lengths on reproduction and neuronal RFRP-3, Kisspeptin, and GnRH in female Turkish Hamsters. J Biol Rhythms. 2014; 29(3):181-191. https://doi.org/10.1177/0748730414532423
54. Li SN, Xue HL, Zhang Q, *et al*. Photoperiod regulates the differential expression of KiSS-1 and GPR54 in various tissues and sexes of striped hamster. Genet Mol Res. 2015; 14(4):13894-13905. http://dx.doi.org/10.4238/2015.October.29.10
55. Klosen KR, Simonneaux V, Klosen P. Differential response patterns of Kisspeptin and RFamide-related peptide to photoperiod and sex steroid feedback in the Djungarian hamster (*Phodopus sungorus*). J Neuroendocrinol. 2017; 29(9):e12529. https://doi.org/10.1111/jne.12529
56. Li Q, Rao A, Pereira A, Clarke IJ, Smith JT. Kisspeptin cells in the ovine arcuate nucleus express prolactin receptor but not melatonin receptor. J Neuroendocrinol. 2011; 23(10):871-882. https://doi.org/10.1111/j.1365-2826.2011.02195.x
57. Gingerich S, Wang X, Lee PKP, *et al*. The generation of an array of clonal, immortalised cell models from the rat hypothalamus: Analysis of melatonin effects on kisspeptin and gonadotropin-inhibitory hormone neurons. Neurosci. 2009; 162(4):1134-1140. https://doi.org/10.1016/j.neuroscience.2009.05.026
58. Dan N, Shah H, Bhatt H, *et al*. Decoding the effect of photoperiodic cues in transducing kisspeptin-melatonin circuit during the pubertal onset in common carp. Mol Reprod Dev. 2024; 91(5):e23744. https://doi.org/10.1002/mrd.23744
59. Zhang Y, Proenca R, Maffei M, Positional cloning of the mouse obese gene and its human homologue. Nature. 1994; 372:425-432. https://doi.org/10.1038/372425a0
60. Gautron L, Elmquist JK. Sixteen years and counting: An update on leptin in energy balance. J Clin Invest. 2011; 121(6):2087-2093. https://doi.org/10.1172/JCI45888
61. Smith JT, Acohido BV, Clifton DK, Steiner RA. KiSS-1 neurones are direct targets for leptin in the *ob/ob* mouse. J Neuroendocrinol. 2006; 18(4):298-303. https://doi.org/10.1111/j.1365-2826.2006.01417.x
62. Donato J, Cravo RM, Frazao R, *et al*. Leptin's effect on puberty in mice is relayed by the ventral pre-mammillary nucleus and does not require signaling in Kiss1 neurons. J Clin Invest. 2011; 121(1):355-368. https://doi.org/10.1172/JCI45106
63. Roa J, Galiano DG, Varela L, *et al*. The mammalian target of rapamycin as novel central regulator of puberty onset *via* modulation of hypothalamic Kiss1 system. Endocrinol. 2009; 150(11):5016-5026. https://doi.org/10.1210/en.2009-0096
64. Luque RM, Kineman RD, Sempere MT. Regulation of hypothalamic expression of KiSS-1 and GPR54 genes by metabolic factors: Analyses using mouse models and a

cell line. Endocrinol. 2007; 148(10):4601-4611. https://doi.org/10.1210/en.2007-0500
65. Wahab F, Ullah F, Chan YM, *et al*. Decrease in hypothalamic Kiss1 and Kiss1r expression: A potential mechanism for fasting-induced suppression of the HPG axis in the adult male Rhesus monkey (*Macaca mulatta*). Horm Metab Res. 2010; 43(02):81-85. https://doi.org/10.1055/s-0030-1269852
66. Varela L, Horvath TL. Leptin and insulin pathways in POMC and AgRP neurons that modulate energy balance and glucose homeostasis. EMBO Rep. 2012; 13(12):1079-1086. https://doi.org/10.1038/embor.2012.174
67. Fu LY, Pol AN. Kisspeptin directly excites anorexigenic proopiomelanocortin neurons but inhibits orexigenic neuropeptide Y cells by an indirect synaptic mechanism. J Neurosci. 2010; 30(30):10205-10219. https://doi.org/10.1523/JNEUROSCI.2098-10.2010
68. Wahab F, Shahab M, Behr R. The involvement of gonadotropin inhibitory hormone and kisspeptin in the metabolic regulation of reproduction. J Endocrinol. 2015; 225(2):R49-R66. https://doi.org/10.1530/JOE-14-0688
69. Higo S, Iijima N, Ozawa H. Characterisation of Kiss1r (Gpr54)- expressing neurones in the arcuate nucleus of the female rat hypothalamus. J Neuroendocrinol. 2017; 29(2). https://doi.org/10.1111/jne.12452
70. Bentefour Y, Bakker J. Kisspeptin signaling and nNOS neurons in the VMHvl modulate lordosis behavior but not mate preference in female mice. Neuropharmacol. 2021; 198:108762. https://doi.org/10.1016/j.neuropharm.2021.108762
71. Jais A, Bruning JC. Arcuate nucleus-dependent regulation of metabolism-pathways to obesity and diabetes mellitus. Endocr Rev. 2021; 43(2):314-328. https://doi.org/10.1210/endrev/bnab025
72. Hu MH, Li XF, McCausland B, *et al*. Relative importance of the arcuate and anteroventral periventricular Kisspeptin neurons in control of puberty and reproductive function in female rats. Endocrinol. 2015; 156(7):2619-2631. https://doi.org/10.1210/en.2014-1655
73. Voigt C, Gahr M, Bennett NC. Differential regulation of Kiss1 gene expression by oestradiol in the hypothalamus of the female Damaraland mole-rat, an induced ovulator. Gen Comp Endocrinol. 2023; 341:114334. https://doi.org/10.1016/j.ygcen.2023.114334
74. Gottsch ML, Cunningham MJ, Smith JT, *et al*. A role for Kisspeptins in the regulation of gonadotropin secretion in the mouse. Endocrinology. 2004; 145(9):4073-4077. https://doi.org/10.1210/en.2004-0431
75. Herbison AE, Theodosis DT. Immunocytochemical identification of oestrogen receptors in preoptic neurones containing calcitonin gene-related peptide in the male and female rat. Neuroendocrinol. 1992; 56(5):761-764. https://doi.org/10.1159/000126304
76. Uenoyama Y, Inoue N, Nakamura S, Tsukamura H. Kisspeptin neurons and estrogen-estrogen receptor α signaling: Unraveling the mystery of steroid feedback system regulating mammalian reproduction. Int J Mol Sci. 2021; 22(17):9229. https://doi.org/10.3390/ijms22179229
77. Terasawa E, Kurian JR, Guerriero KA, *et al*. Recent discoveries on the control of GnRH neurons in nonhuman primates. J Neuroendocrinol. 2010; 22(7):630-638. https://doi.org/10.1111/j.1365-2826.2010.02027.x
78. Smith JT, Popa SM, Clifton DK, *et al*. Kiss1 neurons in the forebrain as central processors for generating the preovulatory luteinizing hormone surge. J Neurosci. 2006; 26(25):6687-6694. https://doi.org/10.1523/JNEUROSCI.1618-06.2006
79. Ajayi AF, Akhigbe RE. Staging of the estrous cycle and induction of estrus in experimental rodents: An update. Fertil Res Pract. 2020; 6(5). https://doi.org/10.1186/s40738-020-00074-3
80. Moore AM, Coolen LM, Porter DT, *et al*. KNDy cells revisited. Endocrinol. 2018; 159(9):3219-3234. https://doi.org/10.1210/en.2018-00389
81. Dubois SL, Wolfe A, Radovick S, *et al*. Estradiol restrains prepubertal gonadotropin secretion in female mice *via* activation of ERα in Kisspeptin neurons. Endocrinol. 2016; 157(4):1546-1554. https://doi.org/10.1210/en.2015-1923
82. Olster DH, Foster DL. Control of gonadotropin secretion in the male during puberty: A decrease in response to steroid inhibitory feedback in the absence of an increase in steroid-independent drive in the sheep. Endocrinol. 1986; 118(6):2225-2234. https://doi.org/10.1210/endo-118-6-2225
83. Lehman MN. Origins of the 'KNDy hypothesis' of GnRH pulse generation. Nat Rev Endocrinol. 2022; 18(9):521. https://doi.org/10.1038/s41574-022-00703-5
84. Constantin S. Targeting KNDy neurons to control GnRH pulses. Curr Opin Pharmacol. 2022; 67:102316. https://doi.org/10.1016/j.coph.2022.102316
85. Lippincott MF, Leon S, Chan YM, *et al*. Hypothalamic reproductive endocrine pulse generator activity independent of Neurokinin B and Dynorphin signaling. J Clin Endocrinol Metab. 2019; 104(10):4304-4318. https://doi.org/10.1210/jc.2019-00146
86. Roa J, Cruz MR, Pino FR, *et al*. Dicer ablation in Kiss1 neurons impairs puberty and fertility preferentially in female mice. Nat Commun. 2022; 13:4663. https://doi.org/10.1038/s41467-022-32347-4
87. Chen J, Minabe S, Munetomo A, *et al*. Kiss1-dependent and independent release of luteinising hormone and testosterone in perinatal male rats. Endocr J. 2022; 69(7):797-807. https://doi.org/10.1507/endocrj.EJ21-0620
88. Dai R, Xu W, Chen W, *et al*. Epigenetic modification of Kiss1 gene expression in the AVPV is essential for female

reproductive aging. Biosci Trends. 2022; 16(5):346-358. https://doi.org/10.5582/bst.2022.01358

89. Ziarniak K, Yang T, Boycott C, *et al*. DNA hypermethylation of Kiss1r promoter and reduction of hepatic Kiss1r in female rats with Type 2 diabetes. Epigenetics. 2022; 17(13):2332-2346. https://doi.org/10.1080/15592294.2022.2119120
90. Diaz MC, Jaramillo AJ, Soto EY, Elosegui PM. Reproductive disease epigenetics. Medical Epigenetics. 2nd Edition. Elsevier eBooks. 2021; 29:309-346. https://doi.org/10.1016/B978-0-12-823928-5.00027-X
91. Kurian JR. Epigenetic regulation of the GnRH and Kiss1 genes. Epigenetics and Human Health ((EHH). Springer. 2016:243-264. https://doi.org/10.1007/978-3-319-24493-8_11
92. Wyatt AK, Zavodna M, Viljoen JL, Stanton JAL, *et al*. Changes in methylation patterns of Kiss1 and Kiss1r gene promoters across puberty. Genet Epi Insights. 2013. https://doi.org/10.4137/GEG.S12897
93. Toro CA, Wright H, Aylwin CF, *et al*. Trithorax dependent changes in chromatin landscape at enhancer and promoter regions drive female puberty. Nat Commun. 2018; 9:57. https://doi.org/10.1038/s41467-017-02512-1
94. Lomniczi A, Loche A, Castellano JM, *et al*. Epigenetic control of female puberty. Nat Neurosci. 2013; 16(3):281-289. https://doi.org/10.1038/nn.3319
95. Oliveira CCV, Fatsini E, Fernandez I, *et al*. Kisspeptin influences the reproductive axis and circulating levels of microRNAs in *Senegalese sole*. Int J Mol Sci. 2020; 21(23):9051. https://doi.org/10.3390/ijms21239051
96. Tsoutsouki J, Abbara A, Dhillo W. Novel therapeutic avenues for kisspeptin. Curr Opin Pharmacol. 2022; 67:102319. https://doi.org/10.1016/j.coph.2022.102319