



# Efficacy of Circuit Training on Cardiorespiratory Fitness, Body Mass Index and Quality of Life in Menopausal Women

Sakeena Patel\* and Chandrakant Patil

Department of Physiotherapy, Krishna Vishwa Vidyapeeth (Deemed to be University), Karad – 415110, Maharashtra, India; patelsakeenaakbar@gmail.com

## Abstract

**Aim:** An adult female population that appeared to be in good health was included in this study to assess the effects of a six-week circuit training programme on cardiovascular endurance and quality of life. **Background:** People going through menopause are looking for ways to improve their quality of life and alleviate symptoms, and hormone medication has a lot of side effects, therefore people are looking towards non-pharmacological approaches. Several studies have shown that aerobic and resistance training have positive effects on women going through menopause. **Method:** The study included 44 menopausal women who were randomly allocated to either a group that trained using circuit training for six weeks or a control group that had regular physiotherapy for the same amount of time. We tracked BMI before and after the intervention, along with cardiorespiratory fitness indicators such as heart rate, oxygen saturation, blood pressure, and the distance covered in a six-minute walk test. Additionally, we administered the Menopause-Specific Quality of Life questionnaire. **Statistical Analysis Used:** We used INSTAT 3.06, a statistical programme, and an unpaired t-test to analyze the data. **Results:** The study found that menopausal women's cardiovascular endurance and quality of life were both greatly enhanced by the six-week circuit training programme. "There was no statistically significant change between the pre- and post-intervention values of the body mass index or the Menopause-Specific Quality of Life scores, however, there was a difference in the cardiorespiratory indicators and physical symptoms in quality of life. **Conclusion:** Circuit training has the potential to alleviate menopausal symptoms and improve the quality of life for women going through menopause. Physical therapists treating menopausal women might think about using circuit training as an intervention.

**Keywords:** Body Mass Index, Cardiorespiratory Fitness, Circuit Training, Menopause, Quality of Life (QOL)

## 1. Introduction

In order to keep the community healthy, it is essential to take care of women's health concerns because they are an integral part of both the family and society. A woman enters a new phase of her life when menopause occurs, the end of her reproductive years. One year following the start of the last menstrual cycle is considered the official start of menopause by the World Health Organisation<sup>1</sup>. During the menopausal transition, which begins with irregular bleeding and continues for an average of 3.8 years, the

menstrual cycle ends after a 12-month irregular period. During menopause, a variety of symptoms manifest as a result of decreased estrogen and progesterone levels. The term "perimenopausal" describes a woman's condition when she is just entering menopause<sup>1,2</sup>. The postmenopausal condition is characterized by the persistence of menopausal symptoms in women beyond menopause<sup>1,2</sup>. A woman's life might be drastically altered by menopause symptoms. In both the perimenopausal and postmenopausal stages, the majority of women (85%)

\*Author for correspondence

report experiencing multiple menopausal symptoms<sup>3</sup>. Physical symptoms include pain in the joints and the need to urinate often; psychosocial symptoms include sadness, mood swings, and poor cognitive function; and sexual symptoms include a dry vagina and a decrease in desire for sex. Hot flashes and sweats are examples of vasomotor symptoms<sup>4</sup>. Several aspects of Quality of Life (QOL) may be diminished when menopausal symptoms are present<sup>5</sup>.

Some women seek out alternative therapies to alleviate menopausal symptoms and improve their overall well-being. The present treatment options include Hormone Therapy (HT) and CAM (Complementary and Alternative Medicine)<sup>5,6</sup>. Alternative and complementary medicine encompasses a wide range of treatments, such as aromatherapy, massage, meditation, and movement awareness programmes. These treatments can supplement or even replace more conventional medical practices<sup>6</sup>. Many women are seeking alternative treatments for problematic symptoms because of the adverse effects of HT<sup>7</sup>. Examining the ways in which exercise alleviates menopausal symptoms is, thus, of paramount importance.

In Circuit Training (CT), you'll use weight training equipment in between different cardio workouts. At each station, you're required to complete a workout for a specific duration. Once all stations have been finished, one circuit will be shown<sup>8,9</sup>. Evidence suggests that aerobic and resistance training can help women in both the pre- and post-menopausal stages of their bodies' life cycles<sup>5,10,11</sup>. CT can have varying degrees of intensity. Increasing cardiovascular endurance is more likely to be achieved through High-Intensity Interval Training (HIIT) rather than moderate-intensity exercise. Maximal oxygen consumption, maximal pulmonary ventilation, functional capacity, myocardial strength, power, and endurance are some of the cardiopulmonary metrics that are enhanced by<sup>12,13</sup> CT which includes endurance workouts, leading to higher cardiovascular endurance<sup>14</sup>. Enhanced hemodynamic measures, including mean arterial pressure, cardiac output, and heart rate, are the result of increased cardiorespiratory fitness<sup>15</sup>.

Consequently, we set out to see whether postmenopausal women could enhance their cardiorespiratory fitness, Body Mass Index (BMI), and Quality of Life by participating in a circuit training programme for six weeks.

## 2. Method

### 2.1 Design

Pre- and post-design experimental investigation. Circuit training's effects on postmenopausal women's cardiorespiratory fitness, BMI, and Quality of Life were the focus of this research.

### 2.2 Sample Size and Randomization

A formula was used to determine the sample size, which is  $n = Z^2 * P * Q / L^2$ . We estimated a 0.5 standard deviation between the prevalence of menopausal symptoms and quality of life in India, and we found that a reasonable sample size of 44 was required for a 95% confidence range. Equal numbers of eligible individuals were assigned to each sample at random.

### 2.3 Subject Criteria

The selection process resulted in 44 participants meeting the inclusion and exclusion criteria. Sedentary middle-aged women (45-65 years old) were the subjects of the study. We excluded subjects who had a history of gynaecological surgery, or systemic diseases, were taking any medication (including antidepressants and antipsychotics), or were undergoing hormone replacement treatment.

### 2.4 Outcome Measures

- i) Body Mass Index: Body Mass Index (BMI) was computed using the formula  $BMI = \text{kg/m}^2$  after measuring height (with a height rod) and weight (using an electronic balance).
- ii) Cardiorespiratory fitness: This term describes how well the respiratory and circulatory systems can provide oxygen to working muscles when they are subjected to prolonged physical exertion. Cardiovascular endurance is another name for it. The protocol's pre- and post-intervention measurements are based on a six-minute walk test. Effects on systolic and diastolic blood pressure, heart rate, oxygen saturation, and respiratory rate were used to estimate cardiorespiratory fitness. Using a stethoscope and sphygmomanometer, blood pressure was measured. In a six-minute walk test, the entire distance covered is called the six-minute walk distance. An oxygen saturation level, heart rate, and respiratory rate were

all measured with the use of a pulse oximeter and a stopwatch, respectively.

- iii) Quality of Life: The MENQOL questionnaire, which is self-administered and has 29 Likert-scale items, is used for this measurement. You might think of the menopause symptoms on this scale as falling into four categories: vasomotor (items 1-3), psychosocial (items 4-10), physical (items 11-26), and sexual (items 27-29). In where 0 signifies “not bothered at all” and 6 “extremely bothered”<sup>4</sup>.

## 2.5 Procedure

After the institutional ethics committee gave their mandate, participants were chosen using predetermined criteria for inclusion and exclusion. The subjects were informed of the purpose and process in their familiar language. Subject rights were safeguarded and the participant’s informed consent was obtained voluntarily. The participant’s demographic information was recorded. People were chosen for the sample according to the inclusion and exclusion criteria.

Each of the two groups (labelled A and B) consisted of 22 people who took part in the research. The participant’s height, weight, cardiorespiratory fitness indices, and MENQOL scores were evaluated before the intervention began. Afterwards, the protocol was put into action according to the groups. People in the study (group A) did circuit training for 50 minutes three times a week for about six weeks, whereas those in the control group (group B) did traditional physiotherapy for 50 minutes until the end of the study. The aforementioned measures were used in the subsequent post-intervention evaluation. After keeping track of the numbers, we compared them between the two sets of data to see whether they were statistically significant.

For Group A: The circuit training programme began with a 5-minute warm-up that included activities like jogging, stretching, or marching in place. Then, participants moved on to a main class that included six workout stations. With a 30-second break in between, each exercise was performed 10-15 times. The activities that made up the high-intensity interval circuit training were squats, lunges, push-ups, belly crunches, and jumping jacks. The 5-minute cooldown was very much like the warm-up.

For Group B: Group B’s control subjects participated in traditional aerobic exercises prescribed by physical

therapists for 50 minutes, three times weekly. These exercises included walking, stretching, and breathing exercises.

## 2.6 Statistical Analysis

The initial process’s Mean and Standard Deviation were computed by statistical analysis in Microsoft Excel. The disparities between the pre- and post-test values were then examined using the statistical software INSTAT, version 3.06. To go a step further with the calculations, we ran an unpaired t-test Welch test.

## 3. Results

Interpretation:

- i) Body Mass Index (BMI): The experimental group’s mean Body Mass Index (BMI) dropped from 27.11 before intervention to 25.3 after, as shown in Table 1. There was no statistically significant relationship between the variables in the unpaired t-test ( $t = 1.594$ ,  $p = 0.1185$ ). There was a small but non-significant drop in body mass index. There was no change in Body Mass Index (BMI) following intervention in the control group (Table 2).
- ii) Cardiorespiratory fitness: Table 1 shows that after the intervention, the experimental group’s mean values of cardiorespiratory fitness indicators (SBP, DBP, 6MWT, HR, SPO<sub>2</sub>, RR) improved. A statistically significant p-value of 0.0161, 0.0002, 0.0389, <0.0001, <0.0001, <0.0001 was produced by the unpaired t-test, which also produced t-values of 2.507, 4.031, 2.132, 4.662, 4.733, and 4.373. The results show that cardiorespiratory fitness improves with circuit training. Table 2 showed that the control group’s mean cardiorespiratory fitness index values fell marginally, while the p-value was insignificant.
- iii) Quality of Life: Table 1 shows that after the intervention, the mean values of the MENQOL score improved across the board, with the only exception of physical symptoms, where the improvement was statistically significant. The t-value for physical symptoms was 4.496 and the p-value was less than 0.0001, indicating statistical significance. The p values for the remaining symptoms, including vasomotor, psychological, and sexual, were more than 0.05, indicating that they were not statistically significant. After 6 weeks, there was no change in

**Table 1.** Comparison of pre- and post-intervention values in the experimental group

Outcome measures	Experimental Group		P	P < 0.05	T
	MeanSD				
	Pre	Post			
BMI	27.113.66	25.33.59	0.1185	No	1.594
Cardiorespiratory Fitness Indices					
SBP	131.65.52	127.25.87	0.0161	Yes	2.507
DBP	854.77	78.954.95	0.0002	Yes	4.031
6MWT	434.129.49	453.429.08	0.0389	Yes	2.132
HR	75.915.79	68.144.98	<0.0001	Yes	4.662
Spo2	97.550.89	98.680.57	<0.0001	Yes	4.733
RR	20.321.39	18.591.15	<0.0001	Yes	4.373
MENQOL					
Vasomotor	3.730.55	3.640.58	0.597	No	0.5327
Psychological	2.950.58	2.910.61	0.8006	No	0.2542
Physical	3.320.57	2.590.5	<0.0001	Yes	4.496
Sexual	2.270.55	2.230.53	0.7813	No	0.2794
Total	12.270.94	11.36	0.0016	Yes	3.378

**Table 2.** Comparison of pre- and post-intervention values in the control group

Outcome measures	Control Group		P	P < 0.05	T
	MeanSD				
	Pre	Post			
BMI	27.513.78	27.513.78	>0.9999	No	0
Cardiorespiratory Fitness Indices					
SBP	1336.68	131.58.16	0.6321	No	0.6321
DBP	84.32 4.07	82.594.16	0.1813	No	1.359
6MWT	431.834.46	43332.6	0.9131	No	0.1098
HR	75.26.27	73.276.65	0.3325	No	0.9804
Spo2	97.450.84	97.680.87	0.3944	No	0.8605
RR	20.231.35	19.911.44	0.4641	No	0.7388
MENQOL					
Vasomotor	3.680.57	3.680.57	>0.9999	No	0
Psychological	2.910.53	2.910.53	>0.9999	No	0
Physical	3.360.58	3.360.58	>0.9999	No	0
Sexual	2.230.53	2.230.53	>0.9999	No	0
Total	12.180.91	12.180.91	>0.9999	No	0

the control group's mean values of the MENQOL score (Table 2).

## 4. Discussion

Undoubtedly, our study's experimental group showed statistically significant improvements in the cardiorespiratory measures, including blood pressure,

oxygen saturation, heart rate, and 6-minute walk test distance. The benefits of circuit exercise programmes for postmenopausal and perimenopausal women have been studied before, but no study has linked circuit training to all of those problems at once. A case-control research was carried out by Miszko and Cress<sup>16</sup> to determine the efficacy of aerobic and resistance training on women experiencing menopause and those just entering the

menopause. Both types of exercise were found to have statistical significance<sup>16</sup>. Asikaine *et al.*,<sup>17</sup> looked at the feasibility and physical advantages of circuit training for menopausal women and found that resistance training improved strength in the lower extremities. According to Kemmler *et al.*,<sup>18</sup> a four-year circuit training programme that tackled the osteoporotic risk of menopause included comprehensive physical assessments. This study provides more evidence that menopausal women can improve their cardiorespiratory fitness with a 6-week circuit training programme. There have been a limited number of research looking at the effects of circuit training on these parameters in postmenopausal women, compared to the number of studies looking at adults. Cardiorespiratory fitness measures have shown substantial gains in other populations that have used circuit training programmes<sup>19-29</sup>. The reduction in body mass index was not statistically significant as we did not prescribe any special diets for the participants in our study.

Furthermore, in terms of physical symptoms, the experimental group in our study showed a slight improvement in quality of life. Mastrangelo *et al.*,<sup>30</sup> found that the experimental group's quality of life significantly improved after participating in circuit training exercises, and Teoman *et al.*,<sup>31</sup> found that menopausal women undergoing hormone therapy who participated in a circuit training programme had a statistically significant improvement in their quality of life.

One out of six female deaths is caused by congestive heart disease, and most of those women aren't active enough to lower their risk<sup>32-34</sup>. This study placed a heavy emphasis on the MENQOL evaluation tool because of the importance of menopause symptoms and Quality of Life (QOL) throughout the menopausal transition. Some of the most common menopausal symptoms include night sweats, mood swings, depression, increased urination, joint discomfort, dry vagina, and decreased desire to have sexual relations. This study highlighted the importance of investigating the impact of this intervention on menopausal symptoms and quality of life as a potential future treatment option.

Improving cardiovascular endurance and quality of life may be possible with a six-week circuit training programme, according to our study's most important findings. In addition, the total training time in this study was just six weeks, which is not a very long time to see a change in body mass index. The lack of longitudinal follow-up and the small sample size were the main

drawbacks of this investigation because of time restrictions. Future studies should incorporate longitudinal follow-up to examine the benefits of circuit training over the long run. To further validate the importance of these findings, an additional study with a bigger sample size is required.

## 5. Conclusion

Cardiorespiratory fitness and quality of life can be enhanced by a brief (6-week) circuit training programme for women going through menopause, according to the current study. Blood pressure, oxygen saturation, heart rate, and 6-minute walk time were all cardiorespiratory indices that showed statistically significant improvement. Although there was no change in the MENQOL scores for vasomotor, psychological, or sexual symptoms after the intervention, there was a significant improvement in physical symptoms. Afterwards, Body Mass Index (BMI) decreased, however the decline was not statistically significant. These results suggest that women going through menopause or just after menopause can benefit from circuit exercise as an intervention. It may take more time and a larger sample size to see whether this intervention makes a significant difference.

**Abbreviations Used:** Circuit Training (CT), Body Mass Index (BMI), Menopause Specific Quality of Life (MENQOL), Blood Pressure (BP), Heart Rate (HR), Oxygen Saturation (SPO<sub>2</sub>), 6 Minute Walk Test Distance (6MWT D)

## 6. Key Points

**Findings:** The circuit training program for menopausal women can improve cardiorespiratory fitness and Quality of Life.

**Implications:** These findings imply that circuit exercise might be effective for menopausal women.

**Limitations:** The study's limitations were the small sample size and the shorter period due to time constraints.

**Caution:** There is a lot of opportunity to obtain important insights and have more influence if this program is used more widely and over a longer length of time.

## 7. References

- Sherman S. Defining the menopausal transition. *Am J Med.* 2005; 118:3-7. <https://doi.org/10.1016/j.amjmed.2005.10.006> PMID:16414321
- Blake J. Menopause: Evidence-based practice. *Best Pract Res Clin Obstet Gynecol.* 2006; 20(6):799-839. <https://doi.org/10.1016/j.bpobgyn.2006.07.001> PMID:17084674
- Woods NE, Mitchell ES. Symptoms during the perimenopause: prevalence, severity, trajectory, and significance in women's lives. *Am J Med.* 2005; 118:14-24. <https://doi.org/10.1016/j.amjmed.2005.10.008>
- Lewis JE, Hilditch JR, Wong CJ. Further psychometric property development of the Menopause-Specific Quality of Life questionnaire and development of modified version MENQOL-Intervention questionnaire. *Maturitas.* 2005; 50:209-221. <https://doi.org/10.1016/j.maturitas.2004.06.015> PMID:15734602
- Teoman N, Ozcan A, Acar B. The effect of exercise on physical fitness and quality of life in postmenopausal women. *Maturitas.* 2004;47(1):71-77. [https://doi.org/10.1016/S0378-5122\(03\)00241-X](https://doi.org/10.1016/S0378-5122(03)00241-X) PMID:14706768
- CAM Basics. National Institutes of Health, National Center for Complementary and Alternative Medicine. Available from: <http://nccam.nih.gov/health/whatiscam/>
- Rossouw JE, Anderson GL, Ross PL, *et al.* Risks and benefits of estrogen plus progestin in healthy postmenopausal women, principal results from the women's health initiative randomized controlled trial. *JAMA.* 2002; 288(3): 321-333. <https://doi.org/10.1001/jama.288.3.321> PMID:12117397
- Jacobs PL, Nash MS, Rusinowski JW. Circuit training provides cardiorespiratory and strength benefits in persons with paraplegia. *Med Sci Sports Exerc.* 2001; 33(5):711-717.
- Maiorana A, O'Driscoll G, Goodman C, Taylor R, Greed D. Combined aerobic and resistance exercise improves glycemic control and fitness in type 2 diabetes. *Diab Res Clin Pract.* 2002; 56:115-123. [https://doi.org/10.1016/S0168-8227\(01\)00368-0](https://doi.org/10.1016/S0168-8227(01)00368-0) PMID:11891019
- Teixeria PJ, Going SB, Houtkooper LB, *et al.* Resistance training in postmenopausal women with and without hormone therapy. *Med Sci Sports Exerc.* 2003; 35(4):555-562.
- Astrand LL, Nedstrand E, Wyon Y, Hammar M. Vasomotor symptoms and quality of life in previously sedentary postmenopausal women randomized to physical activity of estrogen therapy. *Maturitas.* 2004; 48:97-105. [https://doi.org/10.1016/S0378-5122\(03\)00187-7](https://doi.org/10.1016/S0378-5122(03)00187-7) PMID:15172083
- Bocalini DS, Lima LS, de Andrade S, Madureira A, Rica RL, dos Santos RN, Serra AJ, Silva Jr JA, Rodriguez D, Figueira Jr A, Pontes Jr FL. Effects of circuit-based exercise programs on the body composition of elderly obese women. *Clin Interv Aging.* 2012; 7:551. <https://doi.org/10.2147/CIA.S33893> PMID:23271901 PMID:PMC3526879
- Beale L, McIntosh R, Raju P, Lloyd G, Brickley G. A comparison of high intensity interval training with circuit training in a short-term cardiac rehabilitation programme for patients with chronic heart failure. *International Journal of Physical Medicine and Rehabilitation.* 2013; 1:17
- Kumar P. The effect of circuit training on cardiovascular endurance of high school boys. *Global Journal of Human Social Science, Arts, Humanities and Psychology, Diakses.* 2013; 2:13.
- Villelabeitia-Jaureguizar K, Vicente-Campos D, Senen AB, Jiménez VH, Garrido-Lestache ME, Chicharro JL. Effects of high-intensity interval versus continuous exercise training on post-exercise heart rate recovery in coronary heart-disease patients. *Int J Cardiol.* 2017; 244:17-23. <https://doi.org/10.1016/j.ijcard.2017.06.067> PMID:28648356
- Miszko TA, Cress ME. A lifetime of fitness: exercise in the perimenopausal and postmenopausal woman. *Clin Sports Med.* 2000; 19(2):215-232 [https://doi.org/10.1016/S0278-5919\(05\)70200-3](https://doi.org/10.1016/S0278-5919(05)70200-3) PMID:10740756
- Asikainen TM, Suni JH, Pasanen ME, *et al.* Effect of brisk walking in 1 or 2 daily bouts and moderate resistance training on lower extremity muscle through menopause: a randomized, controlled trial. *Phys Ther.* 2006; 86(7): 912-923.
- Kemmler W, Engelke K, Von Stengel S, Weineck J, Lauber D, Kalender WA. Long-term four-year exercise has a positive effect on menopausal risk factors: the Erlangen Fitness Osteoporosis Prevention Study. *J Strength Cond Res.* 2007; 21(1):232-239.
- Villelabeitia-Jaureguizar K, VicenteCampos D, Senen AB, Jiménez VH, Garrido-Lestache ME, Chicharro JL. Effects of high-intensity interval versus continuous exercise training on post exercise heart rate recovery in coronary heart-disease patients. *International Journal of Cardiology.* 2017; 244:17-23. <https://doi.org/10.1016/j.ijcard.2017.06.067> PMID:28648356
- Blair SN, Kohl HW, Paffenbarger RS, Clark DG, Cooper KH, Gibbons LW. Physical fitness and all-cause mortality: A prospective study of healthy men and women. *JAMA.* 1989; 262(17):2395-401. <https://doi.org/10.1001/jama.1989.03430170057028> PMID:2795824
- Booth FW, Laye MJ, Lees SJ, Rector RS, Thyfault JP. Reduced physical activity and risk of chronic disease: The biology behind the consequences. *Eur J Appl Physiol.* 2008; 102(4):381-90. <https://doi.org/10.1007/s00421-007-0606-5> PMID:17987311
- Weston KS, Wisløff U, Coombes JS. High intensity interval training in patients with lifestyle-induced cardio-metabolic

- disease: A systematic review and meta-analysis. *Br J Sports Med.* 2014;48(16):1227-34.
23. Venturelli M, Cè E, Limonta E, Schena F, Caimi B, Carugo S, Veicsteinas A, Esposito F. Effects of endurance, circuit, and relaxing training on cardiovascular risk factors in hypertensive elderly patients. *Age.* 2015;37(5):101. <https://doi.org/10.1007/s11357-015-9835-4> PMID:26381921 PMCid:PMC5005840
  24. Takeshima N, Rogers ME, Islam MM, Yamauchi T, Watanabe E, Okada A. Effect of concurrent aerobic and resistance circuit exercise training on fitness in older adults. *Eur J Appl Physiol.* 2004; 93(1-2):173-182. <https://doi.org/10.1007/s00421-004-1193-3> PMID:15293053
  25. Brentano MA, Cadore EL, Da Silva EM, Ambrosini AB, Coertjens M, Petkowicz R, Viero I, Kruegel LF. Physiological adaptations to strength and circuit training in postmenopausal women with bone loss. *J Strength Cond Res.* 2008; 22(6):1816- 1825. <https://doi.org/10.1519/JSC.0b013e31817ae3f1> PMID:18978624
  26. Romero-Arenas S, Martínez-Pascual M, Alcaraz PE. Impact of resistance circuit training on neuromuscular, cardiorespiratory and body composition adaptations in the elderly. *Aging and Disease.* 2013; 4(5):256.
  27. Ignico AA, Mahon AD. The effects of a physical fitness program on low-fit children. *Research Q Exercise Sport.* 1995; 66:85-90. <https://doi.org/10.1080/02701367.1995.10607659> PMID:7777700
  28. Annesi J, Westcott W, Faigenbaum A, Unruh J. Effects of a 12 week physical activity protocol delivered by YMCA afterschool counsellors (Youth Fit for Life) on fitness and self-efficacy changes in 5-12 year old boys and girls. *Research Q Exercise Sport.* 2005; 76:468-476. <https://doi.org/10.1080/02701367.2005.10599320> PMID:16739685
  29. Wong PCH, Chia MYH, Tsou IYY, Wansaicheong GKL, Tan B, Wang JCK, Tan J, Kim C, Boh G, Lim D. Effects of a 12-week exercise training programme on aerobic fitness, body composition, blood lipids and C-reactive protein in adolescents with obesity. *Ann Acad Med Singapore.* 2008; 37:286-293. <https://doi.org/10.47102/annals-acadmedsg.V37N4p286> PMID:18461212
  30. Mastrangelo MA, MacFarlane S, Woodrow K, Conway E, Klitz DD, Mauriello K, MillerScales A, Nieves-Ventimeglia L, Galantino ML. Effect of circuit training on menopausal symptoms and quality of life. *Journal of Women's Health Physical Therapy.* 2010; 34(2):58-63. <https://doi.org/10.1097/JWH.0b013e3181e8fe09>
  31. Teoman N, Özcan A, Acar B. The effect of exercise on physical fitness and quality of life in postmenopausal women. *Maturitas.* 2004; 47(1):71-77. [https://doi.org/10.1016/S0378-5122\(03\)00241-X](https://doi.org/10.1016/S0378-5122(03)00241-X) PMID:14706768
  32. Lloyd-Jones D, Adams R, Carnethon M, *et al.* Heart disease and stroke statistics2009 update: A report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation.* 2009;119(3):e21-e181.
  33. World Health Organization. Physical activity. Available from: <https://www.who.int/newsroom/factsheets/detail/physical-activity>
  34. Johns NA, Kellar-Guenther Y, Jankowski CM, Neff H, Erlandson KM. A qualitative focus group study of perceived barriers and benefits to exercise by self-described exercise status among older adults living with HIV. *BMJ Open.* 2019; 9(3):e026294. <https://doi.org/10.1136/bmjopen-2018-026294> PMID:30850416 PMCid:PMC6429843