



Research Article

Plasticity in predation behaviour of carabid beetles in agro-ecosystems

N. YAIPHABI CHANU*, R. NAGAR, A. K. MEENA and R. SWAMINATHAN

Department of Entomology, Rajasthan College of Agriculture, Maharana Pratap University of Agriculture and Technology, Udaipur – 313001, Rajasthan, India

*Corresponding author E-mail: naoremyaiphabichanu@yahoo.in

ABSTRACT: A study on the feeding behaviour of three species of *Chlaenius* and two species of *Calosoma* (Coleoptera: Carabidae), the caterpillar and semi-looper hunting predatory ground beetles in agro-ecosystems, was carried out under laboratory conditions in the Department of Entomology, Rajasthan College of Agriculture, MPUAT, Udaipur, during September to November, 2016. The carabid predator species and the second/ third instar tobacco caterpillar prey relationship followed the typical Holling's Type II functional response. Of the three species of *Chlaenius* evaluated, *Chlaenius udaipurensis* Chanu and Swaminathan consumed the maximum prey with a preference for the second instar tobacco caterpillar. Increase in second instar caterpillar prey density from 1 to 5 resulted in an increase in mean prey consumption up to 4.4 in 24 hours. Under enhanced prey provision from 5 to 25 to *Ch. udaipurensis* the maximum mean consumption was 9.0; whereas, the maximum third instar caterpillar consumption was only 2.0 in 24 hours. Both species of *Calosoma* could kill 11 to 13 third instar tobacco caterpillars per day. Increased prey density evinced a gradual decreasing trend in the per cent feeding propensity with little variation among the species for both the genera of carabids. The relationship between prey consumption and the body mass increase for the species of *Chlaenius* showed significant positive correlation ($r = 0.902$ for II instar prey and $r = 0.711$ for III instar prey); similarly, the relationship for *Calosoma* species had significant positive correlation ($r = 0.795$).

KEY WORDS: *Calosoma*, *Chlaenius*, Feeding Behaviour, Functional Response

(Article chronicle: Received: 06-06-2017; Revised: 22-02-2018; Accepted: 03-03-2018)

INTRODUCTION

Insect predators are frequently reckoned as effective regulators of insect pest populations (DeBach and Rosen, 1991; Price, 1997; Obrycki, 1998; Cardoso and Lazzari, 2003; Padmalatha *et al.*, 2003), which has led their increasing use in insect pest management programs (Wiedenmann and Smith, 1997; Riudavets and Castane, 1998). Functional response of a predator is one of the important factors regulating population dynamics of predator-prey systems, and functional response curves can be used to infer basic mechanisms underlying predator-prey interactions, clarify co-evolutionary relationships, and enhance biological control (Houck and Strauss, 1985).

The members of the family Carabidae (Coleoptera) are mostly carnivorous, larvae and adults are nocturnal and hence, less known. These beetles exhibit polyphagy of diverse order (Davies, 1953; 1959 and Thiele, 1977), Scher-

ney (1959) indicated the possibility of carabid beetles being used in the biological control of pests and, the idea that they potentially reduce some pest populations was corroborated by Coaker (1996). Different species of *Calosoma* and *Chlaenius* are reported to be potential biocontrol agents of lepidopteran larvae (Shanower and Ranga, 1990; Rajagopal and Kumar, 1992). *Calosoma sayi* DeJean adults, well documented as predators of larval lepidopteran pests in several row crops (Price and Shepard, 1978, House and All, 1981), also consume live adults of spotted cucumber beetle (*Diabrotica undecimpunctata howardi* Barber), field crickets (*Gryllus* sp.), wireworms (*Melanotus* sp.), southern green stinging bug [(*Nezara viridula* (L.)), annual cicada (*Tibicen* sp.) and a variety of dead arthropods (Young, 1984). Few studies have been conducted on predatory potential and prey preference of different species of *Calosoma* and *Chlaenius* (Kanat and Mol, 2008; Weseloh, 1988; Suenaga and Hamamura, 1998; Katiyar *et al.*, 1976), but no work has been conducted on the functional response and correlation

between prey consumption and increase in body mass of *Calosoma imbricatum* Klug, *Calosoma orientale* Hope, *Chlaenius nitidicollis* Dejean, *Chlaenius distigma* Chaudoir and *Chlaenius udaipurensis* Chanu and Swaminathan, thus, the objective of this study was undertaken under laboratory conditions.

MATERIALS AND METHODS

The experiment was conducted in the Department of Entomology, Rajasthan College of Agriculture, MPUAT, Udaipur, during September to November, 2016. The adults of three species of *Chlaenius*, viz., *Chlaenius nitidicollis* Dejean, *Chlaenius distigma* Chaudoir, *Chlaenius udaipurensis* Chanu and Swaminathan, and two species of *Calosoma*, viz., *Calosoma imbricatum* Klug, *Calosoma orientale* Hope were collected from Ajmer, Bhilwara and Udaipur using light traps. The prey, i.e., *Spodoptera litura* (Fabricius) eggs as well as first instar larvae were collected from field and reared in the laboratory on the soybean leaves.

The adult carabid beetles were kept individually in a dessicator (15cm mouth diameter), filled with moist sand at the base and grass above it to maintain humidity, covered with a muslin cloth and fastened with rubber band. All the three species of *Chlaenius* were provided second as well as third instar larvae at different prey densities, i.e., 1, 2, 3, 4 and 5 larvae per day. For the indigenous species, *Chlaenius udaipurensis* Chanu and Swaminathan, enhanced prey densities of second instar *S. litura* were provided, i.e., 5, 10, 15, 20 and 25 larvae per day because they were available in sufficient numbers for the study being locally abundant. For both the species of *Calosoma*, third instar *S. litura* was provided at prey densities 5, 10, 15, 20 and 25 larvae per day. The experiment on each species was replicated five times.

Observations were recorded on the number of *S. litura* larvae-prey consumed by the adult beetles in each replication after 24h of exposure. The beetles were pre-starved for 24h and weight for each adult beetle was taken before and after provision of prey. The functional response to the prey

Table 1. Functional response parameters for *Chlaenius* species fed on 2nd and 3rd instar *Spodoptera litura* (Fabricius) larvae

| Prey per dessicator (Nos.) (H) | Total prey provided | 2 nd instar larvae as prey | | | | | 3 rd instar larvae as prey | | | | |
|--------------------------------|---------------------|---|---|------------------------|--------|------------------|---|---|------------------------|--------|------------------|
| | | Total prey killed | Average prey killed * (H _a) | Feeding propensity (%) | 1/(HT) | 1/H _a | Total prey killed | Average prey killed * (H _a) | Feeding propensity (%) | 1/(HT) | 1/H _a |
| <i>Ch. nitidicollis</i> | | Linear regression: Y = 0.167 + 0.813x (R ² =0.958) | | | | | Linear regression: Y = 1.084 – 0.074x (R ² =0.047) | | | | |
| 1 | 5 | 5 | 1 | 100.00 | 1 | 1 | 5 | 1 | 100.00 | 1 | 1 |
| 2 | 10 | 10 | 2 | 100.00 | 0.5 | 0.5 | 5 | 1 | 50.00 | 0.5 | 1 |
| 3 | 15 | 10 | 2 | 66.67 | 0.33 | 0.5 | 4 | 0.8 | 26.67 | 0.33 | 1.25 |
| 4 | 20 | 15 | 3 | 75.00 | 0.25 | 0.333 | 5 | 1 | 25.00 | 0.25 | 1 |
| 5 | 25 | 14 | 2.8 | 56.00 | 0.2 | 0.357 | 5 | 1 | 20.00 | 0.2 | 1 |
| <i>Ch. distigma</i> | | Linear regression: Y = 0.099 + 0.877x (R ² =0.973) | | | | | Linear regression: Y = 0.329 + 0.617x (R ² =0.84) | | | | |
| 1 | 5 | 5 | 1 | 100.00 | 1 | 1 | 5 | 1 | 100.00 | 1 | 1 |
| 2 | 10 | 10 | 2 | 100.00 | 0.5 | 0.5 | 10 | 2 | 100.00 | 0.5 | 0.5 |
| 3 | 15 | 15 | 3 | 100.00 | 0.33 | 0.333 | 10 | 2 | 66.67 | 0.33 | 0.5 |
| 4 | 20 | 15 | 3 | 75.00 | 0.25 | 0.333 | 9 | 1.8 | 45.00 | 0.25 | 0.556 |
| 5 | 25 | 15 | 3 | 60.00 | 0.2 | 0.333 | 10 | 2 | 40.00 | 0.2 | 0.5 |
| <i>Ch.udaipurensis</i> (1) | | Linear regression: Y = 0.013 + 0.982x (R ² =0.998) | | | | | Linear regression: Y = 0.305 + 0.645x (R ² =0.877) | | | | |
| 1 | 5 | 5 | 1 | 100.00 | 1 | 1 | 5 | 1 | 100.00 | 1 | 1 |
| 2 | 10 | 10 | 2 | 100.00 | 0.5 | 0.5 | 10 | 2 | 100.00 | 0.5 | 0.5 |
| 3 | 15 | 15 | 3 | 100.00 | 0.33 | 0.333 | 10 | 2 | 66.67 | 0.33 | 0.5 |
| 4 | 20 | 20 | 4 | 100.00 | 0.25 | 0.25 | 10 | 2 | 50.00 | 0.25 | 0.5 |
| 5 | 25 | 22 | 4.4 | 88.00 | 0.2 | 0.227 | 10 | 2 | 40.00 | 0.2 | 0.5 |
| <i>Ch.udaipurensis</i> (2) | | Linear regression: Y = 0.084 + 0.602x (R ² =0.95) | | | | | Not applicable | | | | |
| 5 | 25 | 24 | 4.8 | 96.00 | 0.2 | 0.208 | | | | | |
| 10 | 50 | 36 | 7.2 | 72.00 | 0.1 | 0.139 | | | | | |
| 15 | 75 | 44 | 8.8 | 58.67 | 0.07 | 0.114 | | | | | |
| 20 | 100 | 40 | 8 | 40.00 | 0.05 | 0.125 | | | | | |
| 25 | 125 | 45 | 9 | 36.00 | 0.04 | 0.111 | | | | | |

* Data indicating average prey killed is a mean of 5 replications; R² indicates coefficient of determination value

Table 2. Functional response parameters for *Calosoma* species fed on 3rd instar *Spodoptera litura* (Fabricius) larvae

| Prey per dessorator (Nos.) (H) | Total prey provided | C. imbricatum | | | | | C. orientale | | | | |
|--------------------------------|---------------------|---|---|-------------------------|---------|------------------|---|---|-------------------------|---------|------------------|
| | | Total prey killed | Average prey killed * (H _a) | Feeding propen-sity (%) | 1/ (HT) | 1/H _a | Total prey killed | Average prey killed * (H _a) | Feeding propen-sity (%) | 1/ (HT) | 1/H _a |
| 5 | 25 | 25 | 5 | 100.00 | 0.2 | 0.200 | 25 | 5 | 100.00 | 0.2 | 0.200 |
| 10 | 50 | 46 | 9.2 | 92.00 | 0.1 | 0.109 | 50 | 10 | 100.00 | 0.1 | 0.100 |
| 15 | 75 | 54 | 10.8 | 72.00 | 0.07 | 0.093 | 65 | 13 | 86.67 | 0.07 | 0.077 |
| 20 | 100 | 55 | 11 | 55.00 | 0.05 | 0.091 | 64 | 12.8 | 64.00 | 0.05 | 0.078 |
| 25 | 125 | 55 | 11 | 44.00 | 0.04 | 0.091 | 61 | 12.2 | 48.80 | 0.04 | 0.082 |
| Linear regression | | Y = 0.051 + 0.711 x (R ² =0.951) | | | | | Y = 0.034 + 0.789 x (R ² =0.939) | | | | |

* Data indicating average prey killed is a mean of 5 replications; R² indicates coefficient of determination value

Table 3. Effect of prey consumption on body mass increase in species of *Chlaenius*

| Carabid species | Prey provided () | Mean prey consumed in 24 h | | Mean body mass increase (mg) after feeding on | |
|--|-------------------|----------------------------|-------------------|---|-------------------|
| | | Prey (II instar) | Prey (III instar) | Prey (II instar) | Prey (III instar) |
| <i>Ch. nitidicollis</i> | 1 | 1.0 | 1.0 | 2.0 | 3.0 |
| | 2 | 2.0 | 1.0 | 6.0 | 2.0 |
| | 3 | 2.0 | 0.8 | 5.0 | 3.0 |
| | 4 | 3.0 | 1.0 | 9.0 | 2.0 |
| | 5 | 2.8 | 1.0 | 7.0 | 3.0 |
| <i>Ch. distigma</i> | 1 | 1.0 | 1.0 | 1.0 | 3.0 |
| | 2 | 2.0 | 2.0 | 4.0 | 6.0 |
| | 3 | 3.0 | 2.0 | 6.0 | 5.0 |
| | 4 | 3.0 | 1.8 | 6.0 | 6.0 |
| | 5 | 3.0 | 2.0 | 5.0 | 5.0 |
| <i>Ch. udaipurensis</i> | 1 | 1.0 | 1.0 | 1.8 | 7.0 |
| | 2 | 2.0 | 2.0 | 4.8 | 14.0 |
| | 3 | 3.0 | 2.0 | 6.6 | 15.0 |
| | 4 | 4.0 | 2.0 | 8.0 | 15.0 |
| | 5 | 4.4 | 2.0 | 9.0 | 14.0 |
| S. Em. ± | | | | 0.31 | 0.51 |
| C. D. | | | | 1.17 | 1.91 |
| Correlation between prey consumption and body mass increase: | | | | 0.902** | 0.711* |
| Calculated t-value: | | | | 7.55 | 3.64 |
| <i>Ch. udaipurensis</i> (enhanced prey provision) | 5 | 4.8 | - | 11.0 | - |
| | 10 | 7.2 | - | 15.0 | - |
| | 15 | 8.8 | - | 19.0 | - |
| | 20 | 8.0 | - | 17.0 | - |
| | 25 | 9.0 | - | 17.0 | - |
| S. Em. ± | | | | 0.49 | - |
| C. D. | | | | 1.83 | - |
| Correlation between prey consumption and body mass increase: | | | | 0.958* | |
| Calculated t-value: | | | | 5.81 | |

* indicates coefficient of correlation values significant at p = 0.05; * * indicates significance at p = 0.001

was (1959) $H_a = \frac{a \cdot H \cdot T}{1 + a \cdot H \cdot T_h}$ method described by Holling

Where ‘ H_a ’ is the number of attacked prey, ‘ H ’ is the prey density per unit area, ‘ a ’ is the predator’s search rate, ‘ T_h ’ is the handling time per prey and ‘ T ’ is the total time that is equal to the sum of time spent on searching and time spent on handling.

Holling’s equation was transformed to a linear form (Livdahl and Stiven, 1983) as:

$$H_a = \frac{a \cdot H \cdot T}{1 + a \cdot H \cdot T_h} \quad \longrightarrow \quad \left(\frac{1}{H_a} \right) = \left(\frac{1}{a} \right) \left(\frac{1}{H \cdot T} \right) + \left(\frac{T_h}{T} \right)$$

$y = \alpha \cdot x + \beta$

The experiment was carried out following CRD and the data for increase in body mass after prey consumption for different species of *Chlaenius* and *Calosoma* were analyzed. The correlation between mean prey-consumption and subsequent increase in body mass was also calculated and the test of significance was worked out.

RESULTS AND DISCUSSION

The three species of *Chlaenius* when provided with different densities of second instar tobacco caterpillar as prey (1, 2, 3, 4 and 5) and enhanced prey provided for *Ch. udaipurensis* (5, 10, 15, 20 and 25) the maximum average prey consumed per day by *Ch. nitidicollis* and *Ch. distigma* were 3.0 each, 4.4 by *Ch. udaipurensis* and 9.0 by *Ch. udaipurensis* under enhanced prey provision. For the same three species of *Chlaenius* provided with third instar tobacco caterpillar as prey at prey densities 1, 2, 3, 4 and 5, the maximum average prey consumed per day were 1.0 for *Ch. nitidicollis* and 2.0 each for *Ch. distigma* and *Ch. udaipurensis* (Table 1). For the two species of *Calosoma* when third instar tobacco caterpillar was provided as prey in different densities (5, 10, 15, 20 and 25), the maximum average consumption per day was 11.0 for *C. imbricatum* and 13.0 for *C. orientale* (Table 2).

The carabid predator species of *Chlaenius* and the second/third instar tobacco caterpillar prey relationship followed the typical Holling’s Type II functional response. For all the three species of *Chlaenius*, an increase in caterpillar prey density for both the prey, i.e., second and third instar tobacco caterpillar, resulted in an increase in mean prey consumption initially, but later, leveled off after satiation with little variation among the species (Table 1). All the same, an increase in prey density evinced a gradual decreasing trend in the percent feeding propensity with little

variation for *Ch. nitidicollis* fed on second instar *S. litura*. An increase in second instar prey numbers from 1 to 5 resulted in a decrease in feeding propensity from 100 to 56 per cent, then a slight increase to 75 per cent and decrease to 56 per cent for *Ch. nitidicollis*; the remaining species showed a regular decreasing trend and the corresponding values for *Ch. distigma* were 100 to 60 per cent; while for *Ch. udaipurensis* it was 100 to 88 per cent. When the prey density was enhanced ranging from 5 to 25 second instar caterpillars per day for *Ch. udaipurensis*, the feeding response showed a gradual decrease from 96 to 36 per cent (Table 1). In case of third instar tobacco caterpillar as prey, an increase in prey numbers from 1 to 5 resulted in a decrease in feeding propensity from 100 to 20 per cent for *Ch. nitidicollis*, 100 to 40 per cent for both *Ch. distigma* and *Ch. udaipurensis* (Table 1). This shows that the carabid predator (*Chlaenius*) and the prey (second/ third instar tobacco caterpillar) relationship followed the typical Holling’s Type II functional response. Holling’s functional response when transformed into a linear regression showed an increasing trend in feeding. The linear regression equations species-wise have been presented in the Table (1). Likewise, the predator-prey relationship for the two species of *Calosoma* exhibited a typical Holling’s Type II response. The response transformed into a linear regression showed an increasing trend as depicted by the equations (Table 2). An increase in third instar tobacco caterpillar as prey from 5 to 25 depicted a decreasing feeding tendency from 100 to 44 per cent for *C. imbricatum* and 100 to 48.8 per cent for *C. orientale*.

The relationship between prey consumption with a prey density ranging from 1 to 5 caterpillars per day and the body mass increase for the species of *Chlaenius* and *Calosoma* indicated significant positive correlations. For *Chlaenius*, the correlation coefficient (r) was 0.902 (second instar prey) and 0.711 (third instar prey). Under enhanced prey density ranging from 5 to 25 caterpillars per day for *Chlaenius udaipurensis* Chanu and Swaminathan, a significant positive correlation (r = 0.958) was recorded between mean consumption of second instar tobacco caterpillars and the subsequent increase in body mass (Table 3). In the case of *Calosoma* species, the correlation coefficient (r) was 0.795 (third instar prey) for a prey density ranging from 5 to 25 caterpillars per day (Table 4). The increase in body mass after prey consumption in different species of *Chlaenius* and *Calosoma* also indicate that both the carabid genera happen to be efficient caterpillar hunters (bio-agents) at low prey densities.

Few studies have been reported for the prey preference and predation potential of different species of the genera *Chlaenius* and *Calosoma* and lepidopteran larvae as prey.

Table 4. Effect of prey consumption on body mass increase in species of *Calosoma*

| Carabid species | Prey provided () | Mean prey () consumed in 24 h | Mean body mass increase (mg) after feeding |
|--|----------------------|-----------------------------------|--|
| <i>C. imbricatum</i> | 5 | 5 | 58 |
| | 10 | 9.2 | 88 |
| | 15 | 10.8 | 105 |
| | 20 | 11 | 98 |
| | 25 | 11 | 103 |
| <i>C. orientale</i> | 5 | 5 | 85 |
| | 10 | 10 | 146 |
| | 15 | 13 | 178 |
| | 20 | 12.8 | 180 |
| | 25 | 12.2 | 184 |
| S. Em. ± | | | 2.14 |
| C. D. | | | 8.20 |
| Correlation between prey consumption and body mass increase: | | | 0.795* |
| Calculated t-value: | | | 3.70 |

* indicates coefficient of correlation values significant at $p = 0.05$

Chlaenius panagaeoides (Laferte) consumed 1.4 to 6.9 cow pea aphids per day (Rajagopal and Kumar, 1988). *Chlaenius micans* (Fabricius) and *Chlaenius posticalis* Motschulsky, particularly the larvae of these species were reported to have high consumption rates of diamondback moths per individual (91 and 92 early fourth instars) through the entire larval period (Suenaga and Hamamura, 1998). Under laboratory conditions, a single adult of *Chlaenius viridis* Chaudoir could consume 9-tobacco caterpillar [*Spodoptera litura* (F.)] larvae per day. An equal liking for the larvae of cotton leaf roller (*Sylepta derogata*) as prey was also observed; however the larvae of *Chlaenius viridis* Chaudoir were observed to prey upon the soybean leaf webber, *Lamprosema* sp. in the field (Swaminathan *et al.*, 2001). Kanat and Mol (2008) studied the feeding capacity of *Calosoma sycophanta* L., which is one of the most important predators of Pine Processionary Moth (PPM) larvae and pupae. The adult beetles consumed as many as 7-8 times their body weight per day; one adult beetle consumed 210-280 PPM larvae per year and 840-1120 PPM larvae during its lifetime. A detailed study on the feeding behaviour of *Calosomasa yi* DeJean by Young (2008) evinced that there was a trend observed wherein initial offerings of prepupae and pupae of fall armyworm were eagerly consumed, but later offerings were less consumed but no such trend occurred with offerings of larvae and adult fall armyworm.

Earlier studies on the functional response of carabid beetles are very scanty and none for the species of *Chlaenius* and *Calosoma*. The present study on functional response of the above mentioned two genera is supported by the studies of Dinis *et al.* (2016), though in an indirect way, as the predator-prey relationship for both the predators *Calathus granat-*

ensis (Vuillefroy) and *Pterostichus globosus* (Fabricius) with the prey *Bactrocera oleae* (Rossi) pupae exhibited a type II functional response.

REFERENCES

- Cardoso JT, Lazzari SMN. 2003. Comparative biology of *Cycloneda sanguinea* (Linnaeus, 1763) and *Hippodamia convergens* (Guerin - Meneville, 1842) (Coleoptera, Coccinellidae) focusing on the control of *Cinara* spp. (Hemiptera, Aphididae). *Rev Bras Entomol.* **47**: 443–446. <https://doi.org/10.1590/S0085-56262003000300014>
- Coaker TH. 1966. The effect of soil insecticides on the predators and parasites of the cabbage root fly (*Erioschiabrassicae* Bouche) and on the subsequent damage caused by the pest. *Ann Appl Biol.* **57**: 397–407. <https://doi.org/10.1111/j.1744-7348.1966.tb03833.x>
- Davies MJ. 1953. The contents of the crops of some British carabid beetles. *Entomol Mag.* **89**: 18–23.
- Davies MJ. 1959. A contribution to the ecology of *Notiophilus* and allied genera (Coleoptera: Carabidae). *Entomol Mag.* **95**: 25–28.
- De Bach P, Rosen D. 1991. *Biological Control*, 2nd ed. Cambridge Univ. Press, New York. 325 pp.
- Dinis AM, Pereira JA, Benhadi-Marin J, Santos, SAP. 2016. Feeding preferences and functional responses of *Calathus granatensis* and *Pterostichus globosus* (Coleoptera: Carabidae) on pupae of *Bactrocera oleae* (Diptera:

- Tephritidae). *Bull Entomol Res.* **106**: 701–709. <https://doi.org/10.1017/S0007485316000213.20> PMID:27063655
- Holling CS. 1959. Some characteristics of simple types of predation and parasitism. *Can Entomol.* **91**: 385–398. <https://doi.org/10.4039/Ent91385-7>
- House GJ, All JN. 1981. Carabid beetles in soybean agro-systems. *Environ Entomol.* **10**: 194–196. <https://doi.org/10.1093/ee/10.2.194>
- Kanat M, Mol T. 2008. The effect of *Calosoma sycophanta* L. (Coleoptera: Carabidae) feeding on the Pine Processionary Moth, *Thaumetopoea pityocampa* (Denis & Schiffermukker) (Lepidoptera: Thaumetopoeidae), in the laboratory. *Turk J Zool.* **32**: 367–372.
- Katiyar RR, Misra BP, Upadhyay KD, Prasad N. 1976. Laboratory evaluation of carabid larva, *Chlaenius bioculatus* (Col.: Carabidae) as a predator of lepidopterous pests. *Entomophaga.* **21**(4): 349–351. <https://doi.org/10.1007/BF02371632>
- Livdahl TP, Stiven AE. 1983. Statistical difficulties in the analysis of predator functional response data. *Can Entomol.* **115**: 1365–1370. <https://doi.org/10.4039/Ent1151365-10>
- Obrycki JJ. 1998. Predaceous coccinellidae in biological control. *Annu Rev Entomol.* **43**: 295–321. <https://doi.org/10.1146/annurev.ento.43.1.295> PMID:15012392
- Padmalatha C, Singh AJAR, Jeyapauld C. 2003. Predatory potential of syrphid predators on banana aphid, *Pentalonia nigronervosa* Coq. *J Appl Zool.* **14**: 140–143.
- Price JF, Shepard M. 1978. *Calosoma sayi* and *Labidura riparia* predation of noctuid prey in soybeans and locomotor activity. *Environ Entomol.* **7**: 653–656. <https://doi.org/10.1093/ee/7.5.653>
- Price PW. 1997. *Insect Ecology*. 3rd ed. Wiley. 874 pp.
- Rajagopal D, Kumar P. 1992. Carabids (Coleoptera: Carabidae) as potential predators on major crop pests in South India. *J Biol Control.* **6**(1): 13–17.
- Rajgopal D, Kumar P. 1988. Predation potentiality of *Chlaenius panagaeoides* (Laferte) (Coleoptera: Carabidae) on cowpea aphid *Aphis craccivora* Koch (Homoptera: Aphididae). *J Aphidology.* **2**: 93–99.
- Riudavents J, Castane C. 1998. Identification and evaluation of native predators of *Frankliniella occidentalis* (Thysanoptera: Thripidae) in the Mediterranean. *Environ Entomol.* **27**: 86–93. <https://doi.org/10.1093/ee/27.1.86>
- Scherney F. 1959. Unsere Laufkafer – Neme Brehm – Bucherei H. 245 Wittenberg.
- Shanower TG and Ranga GVR. 1990. *Chlaenius* sp. (Col.: Carabidae): A predator of groundnut leaf miner larvae. *Int Arachis Newsl.* (8): 19–20.
- Suenaga H, Hamamura T. 1998. Laboratory evaluation of carabid beetles (Coleoptera: Carabidae) as predators of diamondback moth (Lepidoptera: Plutellidae) larvae. *Environ Entomol.* **27**(3): 767–772. <https://doi.org/10.1093/ee/27.3.767>
- Swaminathan R, Bhati KK, Hussain T. 2001. Preliminary investigations on the predation potential of carabids. *Indian J Appl Entomol.* **15**: 37–41.
- Thiele HU. 1977. *Carabid beetles in their environments*. Springer-Verlag, Berlin. <https://doi.org/10.1007/978-3-642-81154-8>
- Weseloh RM. 1988. Prey preferences of *Calosoma sycophanta* L. (Coleoptera: Carabidae) larvae and relationship of prey consumption to predator size. *Can Entomol.* **120**(10): 873–880. <https://doi.org/10.4039/Ent120873-10>
- Wiedenmann RN, Smith JW. 1997. Attributes of the natural enemies in ephemeral crop habitats. *Biol Control.* **10**: 16–22. <https://doi.org/10.1006/bcon.1997.0544>
- Young OP. 2008. Body weight and survival of *Calosoma sayi* (Coleoptera: Carabidae) during laboratory feeding regimes. *Ann Entomol Soc Am.* **101**(1): 104–112. [https://doi.org/10.1603/0013-8746\(2008\)101\[104:BWASOC\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2008)101[104:BWASOC]2.0.CO;2)