



## Research Article

# Molecular mechanisms adopted by abiotic stress tolerant *Pseudomonas fluorescens* (NBAIL-PFDWD) in response to in vitro osmotic stress

**KARKERA ASHWITHA\*, RAJAGOPAL RANGESHWARAN and GOPALSAMY SIVAKUMAR**

ICAR, National Bureau of Agricultural Insect Resources, Hebbal, Bellary Road, Bangalore – 560024, Karnataka, India

\*Corresponding author E-mail: ashwi1986@gmail.com

**ABSTRACT:** Water stress is one of the limiting factors which influences plant growth. Microbes being a partner are an integral part of the ecosystem which influences plant growth under stress. In the present study, *Pseudomonas fluorescens* (NBAIL-PFDWD) subjected to osmotic stress by altering osmotic potential (-10.28 MPa and -26.82 MPa) using Polyethylene Glycol (PEG) 6000 in its growth media revealed expression of proteins which modulate its cell processes. MALDI TOF studies of selected spots from 2D gel analysis of *P. fluorescens* (NBAIL-PFDWD) grown under different osmotic stresses revealed that the stress kindled genes which were involved in production of osmoprotectants, genes encoding DNA damage repair and increased the translational accuracy. The studies also showed that *P. fluorescens* possesses unique mechanisms for survival under osmotic stress. The studies indicate the diverse expression of proteins in *P. fluorescens* (NBAIL-PFDWD) under different osmotic potentials which helped them to mitigate the impact of osmotic stress. The present method unraveled the mechanisms adopted by *P. fluorescens* (NBAIL-PFDWD) to thrive under osmotic stress. The bacterium is a potential stress-tolerant isolate which can be exploited as a plant growth-promoting rhizobacteria for agricultural crops grown in stressed soils.

**KEY WORDS:** MALDI-TOF, osmotic potential, osmotic stress, *Pseudomonas fluorescens*, PEG 6000

(Article chronicle: Received: 02-12-2017; Revised: 26-02-2018; Accepted: 18-03-2018)

## INTRODUCTION

Bacteria surviving in soil are continuously exposed to a number of stresses *viz.*, flooding, high and low temperatures, drought and chemicals. Scientific studies to understand the mechanisms followed by bacteria are interesting and bring forward the survival strategy ratified by them. Bacteria alter their gene expression patterns producing proteins which help them to cope with these stresses (Boor, 2006). Stress-induced alteration in metabolic pathways including translation is one of the multitude ways to survive under stressful conditions. Protein synthesis is a vital part of cell function as it maintains the fidelity of genetic information (Karas, 2015).

The lipid membrane of Gram-negative bacteria maintains stability by balancing its solute content and the osmotic pressure of the environment. *Pseudomonas* spp. managed the variation in homeostasis by accumulating compatible solutes by *de novo* synthesis, trehalose, glutamate, man-

nitol, glucosylglycerol, ectoine, and hydroxyectoine; other osmotolerance mechanisms include accumulation of K<sup>+</sup> ions, proline, gamma-aminobutyrate, -aminobutyrate, alanine, the quaternary amines glycinebetaine, fully N-methylated amino acid derivatives, sucrose, trehalose (-D-glucopyranosyl-β-D-glucopyranoside) and glucosylglycerol (Kets *et al.*, 1996; Freeman *et al.*, 2010, 2013). Transcriptional regulation of certain genes like ProU, OmpF, OmpC also occurs under osmotic changes (Csonka, 1989). Bacterial sigma factors support bacterial growth and regulate physiological development under stress (Boor, 2006). In addition, membrane transporters and porins facilitate translational stress response under osmotic stress (Wood, 2015). Osmotic potential is used to measure the osmotic property of the bacterial environment.

*Pseudomonas* spp. is ubiquitous and flourishes in diverse habitats. They produce a wide variety of stress response systems to adapt to environmental changes (Udaonda *et al.*, 2012). *Pseudomonas* spp. confers drought resistance by

accumulation of exopolysaccharides and osmolytes which make it notable to use as efficient PGPR in soils (Sandhya *et al.*, 2009).

*Pseudomonas fluorescens* is widely used as a biological control agent but the mechanism it opts to survive under low moisture conditions is less understood. In the present investigation, our objective was to understand mechanisms executed by *Pseudomonas fluorescens* (NBAIL- PFDWD) to survive under osmotic imbalance. *Pseudomonas fluorescens* (NBAIL- PFDWD) isolated from rhizosphere soil is tolerant to high temperature, salinity, pH and osmotic stress (Ashwitha *et al.*, 2013). *In vitro* studies were designed to grow the culture in PEG 6000 mediated osmotic stress in media. PEG, is a polymeric co solvent results in osmotic shifts which lead to change in membrane structure by regulating efflux or influx of solvents to maintain balance (Wood, 1999). *Pseudomonas fluorescens* (NBAIL-PFDWD,

GenBank HM439956) was selected as an abiotic stress tolerant organism during screening of several isolates. It was found tolerant to 45°C, salinity (1.5M NaCl), pH 9.0 and also moisture stress. The protein profiling of the *P. fluorescens* (NBAIL- PFDWD) grown under stress condition was done using 2D gel electrophoresis and major spots corresponding to stress genes were identified using MALDI-TOF analysis.


## MATERIAL AND METHODS

### Osmotic stress mediated growth of *Pseudomonas fluorescens* (NBAIL- PFDWD)

*Pseudomonas fluorescens* (NBAIL- PFDWD GenBank HM439956) was used in the study for investigating the growth under osmotic stress. 50 µl of mid-log-phase culture (12 h) of *P. fluorescens* (NBAIL- PFDWD) was inoculated to 250 ml Erlenmeyer flask containing 100 ml of nutrient broth (g/L<sup>-1</sup>) Peptone, 5g; Beef extract, 3g; NaCl, 5g)

**Table 1. List of selected MALDI- TOF identified proteins expressed by *Pseudomonas fluorescens* (NBAIL-PFDWD) under different osmotic potentials**

Stress condition	Spot no.	Protein name (Swiss Prot Database)	Log Normalized Values		
			0 MPa	-10.28 MPa	-26.82 MPa
Normal growth condition	241	2-isopropylmalate synthase OS	5.34	5.11	4.82
	240	Probable dual-specificity RNA methyltransferase RImN OS	4.95	4.64	4.46
	477	30s ribosomal protein S8 OS	5.54	5.12	5.11
	482	Exodeoxyribonuclease 7 small subunit OS	5.62	5.19	5.05
	336	Bifunctional pantoate ligase/cytidylate kinase OS	4.18	3.94	3.78
	391	Cellular nucleic acid-binding protein homolog OS	4.84	4.34	4.36
	392	Threonine--tRNA ligase OS	4.72	4.48	4.40
30% PEG stress (-10.28 OP)	496	Dihydrorotate dehydrogenase (quinone) OS	5.43	5.54	5.56
	90	Ribosome-recycling factor OS	2.95	2.61	3.18
	147	L-ectoine synthase	4.01	4.44	4.00
	156	Arginine deiminase OS	5.49	5.50	5.48
	222	N-acetylmannosamine kinase OS	4.88	4.91	4.96
	347	DNA replication and repair protein RecF OS	4.56	4.53	4.46
	354	Phosphate acyltransferase OS	5.08	4.99	4.98
367	DNA repair protein RecO OS	5.14	5.01	4.95	
50% PEG stress (-26.82 OP)	429	Transcription elongation factor GreA OS	4.36	4.72	5.01
	426	Elongation factor Ts OS	5.16	5.17	5.23
	486	Crossover junction endodeoxyribonuclease RuvC OS	4.87	4.92	4.93
	184	Pantothenate synthetase OS	4.08	4.31	4.43
	218	60 kDa chaperonin OS	4.20	4.40	4.43
	206	DNA mismatch repair protein MutL OS	4.97	4.73	4.98
	248	V-type proton ATPase subunit E OS	4.50	4.13	4.32
	263	Gamma-glutamyl phosphate reductase OS	4.77	4.81	4.89
	439	Ribonuclease HIII OS	4.87	4.85	4.83
	326	Transcriptional repressor NrdR OS	4.75	4.84	4.94
	355	Leu operon leader peptide OS	4.28	4.44	4.53
	373	Fructose-1,6-bisphosphatase class 2 OS	3.96	4.23	4.37
	396	Putative ATP-dependent Clp protease proteolytic subunit-like OS	4.60	4.47	4.56
	390	Bifunctional protein GImU OS	4.08	4.34	4.44
368	NAD(P)H-quinone oxidoreductase subunit O OS	4.51	4.67	4.68	



**Fig. 1.** Ezzee blue stained SDS PAGE gel after IEF showing different protein spots expressed by *Pseudomonas fluorescens* (NBAII-PFDWD) at different osmotic potentials, A; 0 MPa, B; -10.26 MPa and C; -26.82 MPa. The spots selected for MALDI-TOF are circled in black.

amended with PEG 6000 to attain different osmotic potentials ( ) of 0 MPa, -10.28 MPa and -26.82 MPa concentrations. The pH of the media was adjusted to  $7.2 \pm 2$ . The cultures were incubated at  $28 \pm 2$  °C under constant shaking.

#### Extraction of bacterial proteins

The bacterial cells were collected by centrifugation (10,000 g for 10 min) and washed three times with isotonic solution. The proteins were precipitated according to method described by Paul *et al.* (2006). The protein extracts were frozen until further use.

#### 2D Gel analysis for stress-related proteins

The first dimension was carried out at Genei™ Merck Biosciences, Bangalore, India as per standardized protocol. Gel mixture was prepared containing 10.8 M urea, 30% acrylamide, 20 µl of 100 % Triton-X, 130 µl ampholyte pH (3–10), 3 µl TEMED and APS (10%). The 1D was carried out in EttanIPGphor3 (GE Health care). The gel mixture was poured in tube by using (18-gauge needle) syringe up to 15 cm without any air bubble (remaining 1 cm for sample loading space). Once the gel was set the casted gel tube was fixed to the tank with rubber grommet very tightly (sample loading space of tube facing toward the top reservoir). The reservoir top was filled with 1X cathode buffer (400 mM NaOH) and bottom reservoir with 1X anode buffer (100 mM H<sub>3</sub>PO<sub>4</sub>). The gel was pre-focused by adding 1–3 µl of 2D SLB [9.8 M urea, 2% NP-40 (10% in distilled water), 2% carrier ampholytes 8/10, 25 mM DTT] into sample loading space and power cords were connected to the power supply. Electrophoretic conditions of the rod gels during the IEF (pre-run) were 200 V for 15 min, 300 V for 30 min, and 400 V for 1 h. After IEF (pre-run), 350 µg of sample was loaded and gel was run at 400 V for 16 h. After 16 h, the gels were transferred to a small tray contain-

ing 2D equilibration buffer [6 M urea, 75 mM Tris–HCl (pH 8.8), 29.3% glycerol, 2% SDS, 0.002% bromophenol blue] and left for 30 min.

The second dimension was carried out on 12.5% SDS PAGE. The equilibrated gel was placed on 1% agarose added on the top of the polymerized SDS-gel. Electrophoresis was carried out at 150 V in Genei™ Vertical Maxi Gel system. Protein spots were visualized using Ezzee blue gel stainer (Genei™) and the well-resolved spots were numbered. The Mr of individual protein spots was determined by comparison with molecular weight markers. Spot detection was carried out using the software, Progenesis (Non-linear Dynamics available at [www.nonlinear.com](http://www.nonlinear.com)). The two-dimensional gels of *P. fluorescens* (NBAII- PFDWD) grown in all three osmotic conditions were compared using this software.

#### Protein identification

Selected proteins were subjected to in-gel proteolytic digestion by excising 1-mm-thick gel and washing for 30 min at room temperature under vigorous shaking with 400 µl of 10 mM ammonium bicarbonate solution containing 50% (v/v) acetonitrile. The gel pieces were dried for 15 min in a vacuum concentrator. The rehydrated gel pieces were incubated in 150 µl reduction solution (10 mM DTT, 100 mM ammonium bicarbonate) for 30 min at 56°C. The reduction solution was then discarded and 100 µl alkylation solution (50 mM iodoacetamide, 100 mM ammonium bicarbonate) was added for 30 min in dark at room temperature. The gel pieces were digested in 5 µl trypsin solution (sequencing grade modified trypsin, Promega, Madison and 10 ng/µl in 5 mM ammonium bicarbonate/5% acetonitrile). The samples were incubated for 5 h at 37°C. The reaction was stopped by adding 1 µl of 1 % trifluoroacetic acid. For better extraction of peptides, the samples were stored overnight at 5 °C.

### MALDI-TOF MS Identification of Protein

The analysis was carried out in MALDI-TOF mass spectrometer (Bruker-Daltonics, Germany) at Merck Biosciences, Bangalore, India. The 1 µl of concentrated peptides from the supernatant was mixed with 2 µl of matrix solution [5 mg *o*-cyano-4-hydroxycinnamic acid in 40 % (v/v) acetone, 50 % (v/v) acetonitrile, 9.9 % (v/v) water, and 0.1 % (w/v) TFA in water]. From this mix, 1 µl was deposited onto the MALDI target. Tryptic peptides were analyzed in positive mode. Background ions from trypsin autolysis and contamination by keratins were removed from mass lists. The proteins were analysed using MASCOT search engine in latest version of the NCBI database with following relevant search parameters viz., monoisotopic mass accuracy, peptide mass tolerance (0.1 Da), peptide charge state, allowed variable modifications, oxidation of methionine and fixed modification, carbamidomethylation of cysteine. Fragmentation of selected peptides was measured using the PSD mode. The percentage similarity of amino acids, Mr and pI were taken into consideration for identification of the proteins from bacteria of related species/genera.

### RESULTS AND DISCUSSION


Capability of *P. fluorescens* (NBAIL-PFDWD) to survive osmotic stress is well addressed in the protein profiling technique. The stress response of *P. fluorescens* (NBAIL-PFDWD) subjected to osmotic stress at different osmotic potential using PEG 6000 revealed wide array of proteins produced by translational modification. Overall more than 300 proteins were expressed in all the three growth conditions. Thirty distinctive spots were selected based on Pro-linear software from protein profile gels of all the three growth conditions subjected viz., 0 MPa, -10.28 MPa and -26.82 MPa (Fig. 1) and identified using MALDI-TOF MS. The proteins were abundant under osmotic stress conditions, -10.28 MPa and -26.82 MPa when compared to the normal conditions. Seven spots were detected from protein gel of 0 MPa which were down regulated under osmotic stress (Fig. 1). Eight proteins and 15 proteins which were up regulated under -10.28 MPa and -26.82 MPa were selected for identification respectively. The fold increase of all these proteins are tabulated (Table 1). Comparison of proteins revealed that the expression levels of bifunctional pantoate ligase/cytidylate kinase OS decreased under -10.28 MPa and -26.82 MPa induced osmotic stress. L-ectoine synthase expression was higher under -10.28 MPa induced osmotic stress, however, its expression was almost on par under normal conditions and -26.82 MPa induced stress.

The spot details obtained from Progenesis software clearly distinguishes spots corresponding to transcription

elongation factor GreA OS and fructose-1,6-bisphosphatase class 2 OS were up regulated under both cases of osmotic stress. Their expression under normal condition was barely visible in gel. In both osmotic stress conditions, exodeoxyribonuclease 7 small subunit OS was down regulated. Comparison among all the three gels showed that, arginine deaminase OS, DNA replication and repair protein RecF OS, phosphate acyltransferase OS and DNA repair protein RecO OS showed no significant difference in their expression under normal and osmotic stress conditions (Fig. 2, 3). It was noticed that the 11 proteins selected from both osmotic stress conditions were on par except four identified as transcription elongation factor GreA OS, 60 kDa chaperonin OS, DNA mismatch repair protein MutL OS and Gamma-glutamyl phosphate reductase OS which showed slight increase under -26.82 MPa induced osmotic stress.

In order to survive under moisture stress, *Pseudomonas fluorescens* (NBAIL-PFDWD) produced an array of proteins which helped them mitigate the stress conditions. Most of the proteins shown to over express under moisture condition by *P. fluorescens* (NBAIL-PFDWD) are not reported so far. The current study is novel and has provided an initial platform to explore microbial interaction under stressed ecosystem. Literature studies reveal that all these genes are expressed as stress response in different microorganisms. 2-isopropylmalate synthase was expressed in osmotic stress subjected in *Campylobacter jejuni* and it belongs to transferases involved in L-leucine and pyruvate metabolism (Cameron *et al.*, 2012). RlmN is associated with catalysis of m<sup>2</sup>A synthesis with both rRNA and tRNA. The translation accuracy decreased with inactivation of RlmN in *E. coli* (Benitez-Paez and Villarrayo, 2012). This function reduces the chances of misreading of translational reading frame and further balance bacterial phenotype which could impair under stress conditions. Many genes are expressed for bacterial growth under a stable environment however, its expression varies when the bacteria challenges a distressed milieu. Ribosome profiling provides an insight to its role in expression of bacterial proteins under environmental fluctuations (Starosta *et al.*, 2014). Expression of 30s ribosomal proteins were found to be down regulated in carnocyclin treated *Listeria monocytogenes* (Liu *et al.*, 2014). DNA repair mechanisms play a pivotal role in altering stressed genes for a normal functioning of a cell. Exodeoxyribonuclease 7 small subunit plays important role in DNA repair by avoiding mutation occurring under environment instability (Lau *et al.*, 2011; Lovett, 2011). DNA binding protein modulate biochemical functions, DNA binding and ferroxidase activity to protect cells from damage due to various stresses when in exponential growth and majorly when they enter stationary phase (Karas *et al.*, 2015). No


functional activity was reported for bifunctional pantoate ligase/cytidylate kinase OS under stress conditions. Hence precise regulation of these proteins is important under adverse conditions.



**Fig. 2. A, Spots which are expressed under normal growth conditions (c1) analysed using Progenesis Linear Dynamics Software. Spots 241-392 are shown to be down regulated under conditions c2 (-10.28MPa) and c3 (-26.82 MPa). B, differentially expressed genes under c2.**

Stress involved proteins expression levels switch either by down regulating or up regulating. Eight spots were selected from gel profile of -10.28 MPa induced osmotic stress and fifteen from -26.82 MPa induced osmotic stress, all of which were differentially regulated under stress (Fig. 2, 3). Dihydrorotate dehydrogenase (quinone) OS and arginine deiminase OS were expressed in higher folds under stress and dihydrorotate dehydrogenase (quinone) is involved in pyrimidine biosynthesis however, no function in stress response was observed (Jensen and Larsen, 2003; Kawasaki *et al.*, 2009). Arginine deaminase stimulates the growth of *Pseudomonas* spp. and reverses the adverse effects in acidic environment by serving as energy source (Marquis *et al.*, 1987). Ammonia derived from arginine deaminase breakdown reduced the effect of pH stress in *Staphylococcus epidermidis* (Lindgren *et al.*, 2014). Ribosome-recycling factor OS, L-ectoine synthase, N-acetylmannosamine kinase OS and phosphate acyltransferase OS were over expressed under this stress. Ribosome recycling is closely associated with protein synthesis which is required for bacterial growth. In *E. coli* it was shown that this gene is involve in dissociation of ribosome from mRNA after translation and in turn facilitates next cycle of protein synthesis (Borovinskaya *et al.*, 2007; Li *et al.*, 2012). Increase or decrease in cytoplasmic solutes is primary response to osmotic swing on bacteria. Ectoines belong to an array of compatible solutes produced majorly by halophilic bacteria and are in-

involved in protein stabilization to prevent cell damage under osmotic or temperature stresses (Louis and Galinski, 1997; Kuhlmann *et al.*, 2011). In addition ectoines serve as carbon and nitrogen source or as intracellular reserves (Pastor *et al.*, 2010). Enzymes catalyse metabolic reaction in cell biosynthesis, N-acetylmannosamine kinases catalyse cell biosynthesis and cellular response to DNA damages stimulus. No reports are found with respect to expression of these stress proteins in *P. fluorescens* (NBAIL-PFDWD). Acyltransferase utilizes fatty acid to form membrane phospholipids and further helps in membrane expansion. They regulate membrane biogenesis and cell growth. Notable function under stress is not known (Zhang and Rock, 2008). DNA replication is realm of bacterial survival. Osmotic stress induces perturbations in replication cycle. RecF influence the repair mechanism in responsible damage and replication fork stress in *E. coli* and *B. subtilis* (Lenhart *et al.*, 2014; Villarrayo *et al.*, 1998). Bacteria regulate membrane biogenesis of coordinate cell growth by synthesis of fatty acid. RecO are recombinant mediator protein involved in restarting of stalled replication in ssDNA gaps repair and in repairing dsDNA breaks. It is assisted in rapid repair of chromosome breaks in *Deinococcus radiodurans* under radiation and desiccation stress (Lusetti and Cox, 2002; Ryzhikov *et al.*, 2011). Surprisingly, in the present study DNA replication and repair protein RecF OS and DNA repair protein RecO OS were found to be repressed under osmotic stress.



**Fig. 3. Differentially expressed genes as analysed using Progenesis Linear software under c3 (-26.82 MPa).**

A slight variation in osmotic potential of media prompted expression of proteins to express at higher folds. Nine proteins were up regulated when only 50% water availability was provided in media. GreA cleaves complexes formed at site of transcriptional arrest and resumes the transcriptional process additionally and they are up regulated in environmen-

tal perturbations (Kusuya *et al.*, 2011; Li *et al.*, 2012). The genes which were up regulated were transcription elongation factor GreA OS, elongation factor Ts OS, crossover junction endodeoxyribonuclease RuvC OS, crossover junction endodeoxyribonuclease RuvC OS, pantothenate synthetase OS, 60 kDa chaperonin OS, Gamma-glutamyl phosphate reductase OS, transcriptional repressor NrdR OS, Leu operon leader peptide OS, fructose-1,6-bisphosphatase class 2 OS, bifunctional protein GlmU OS and NAD(P)H-quinone oxidoreductase subunit O OS. RuV is involved in repair of DNA facing damage due to mutations. They encode endonucleases activities which resolve Holliday structure during recombination (Takahagi *et al.*, 1991). No significant function was reported in stress mechanism however its major function lies in protein biosynthesis. 60 kDa chaperonin or commonly referred *E. coli* GroEL is generally exhibited under temperature stress and is involved in unfolding the misfolded protein to its natural state (Maguire *et al.*, 2002; Mande and Kumar, 2013). Gamma glutamyl phosphate reductase catalyses metabolic reaction of a major osmoprotectant proline encoded by ProA. Artificial bifunctional enzyme gamma-GK/gamma GPR expression in *E. coli* increased production of proline and osmotolerance of host (Chen *et al.*, 2006; Liang *et al.*, 2013). It is also associated with production of small amount of glutathione which is linked to proline biosynthesis under oxidative stress (Perez-Arellano *et al.*, 2010). NrdR controls the transcription of genes by binding to promoter region. This regulatory protein controls expression of ribonucleoside reductases which are precursors of DNA synthesis and repair during vegetative growth. These are important for aerobic growth of *E. coli* (Torrents *et al.*, 2007). Fructose 1, 6-bisphosphate class 2 OS play central function in glycolysis. Macomber *et al.* (2011) reported that the cellular over expression of fructose 1,6-bisphosphate in *E. coli* reduced inhibitory effects of nickel. GlmU is a precursor for bacterial cell wall synthesis components like lipopolysaccharides and peptidoglycan. Inhibition of GlmU adversely affects cell viability (Larsen *et al.*, 2012). Hence, this has been utilized as target for antibiotic drug discovery (Mochalkin *et al.*, 2007). EF-Ts serves as the guanine nucleotide exchange factor for EF-Tu which is associated in response to nutrient deprivation and protect from stress (Caldas *et al.*, 1998). NADPH- quinone oxidoreductase subunit is induced to protect cell from oxidative stress along with many other detoxifying enzymes (Atia *et al.*, 2014). In *E. coli* they adapt and overcome a strong imbalance in NADPH redox state (Auriol *et al.*, 2011). Down regulation of three proteins (V-type proton ATPase, Ribosome nuclease II, and Putative ATP-dependent Clp protease proteolytic subunit-like OS) at -26.82 MPa osmotic potential was observed in our study. V-type proton ATPase subunit is a membrane subunit whose major role is not much elucidated. Their function ranges from protein sorting to membrane trafficking. They drive the active transport systems in mem-

brane. These V-type ATPases restrict the influx of external solutes so as to reduce risk of membrane integrity loss (Beyenback and Wiezoerek, 2006; Cohen, 2014). Ribosome nuclease II functions as exoribonucleolytic which is important for cell response to stress conditions. In *E. coli* activity of ribonuclease II was reduced under conditions of slow growth (Song *et al.*, 2016). Similar case was observed in *P. fluorescens* (NBAIL- PFDWD). RNA polymerases transcribe few genes to up regulate when bacteria reach its stationary phase and when its regular process shut downs (Lau *et al.*, 2011). Clp mediates proteolysis of misfolded proteins subsequent to translation (Schelin *et al.*, 2002). They exhibit adaptive response in *B. subtilis* against stationary phase phenomena like sporulation (Msadek *et al.*, 1998). DNA repair was active through other set of genes like MutL. MutL belonging to mismatch repair system suppress chromosomal aberration by correcting mismatch bases during replication error occurring under various stress conditions (Harris *et al.*, 1997) and helps in maintaining genetic stability (Oliver *et al.*, 2002). Unlike other methyl directed mismatch proteins, MutL was functional even in stationary phase of *E. Coli* (Feng *et al.*, 1996). In studies using *P. syringae* it was shown that osmotolerance was influenced by synthesis of N-acetylglutaminylglutamine amide (NAGGN) that causes reduction of cellular glutamine indicating lower glutamine synthetase activity thereby causing glutamate accumulation (Freeman *et al.*, 2013). In our studies up regulation of Gamma-glutamyl phosphate reductase OS was seen and this protein is involved in synthesizes of L-glutamate 5-semialdehyde from L-glutamate. Literature on stress response proteins expressed by a PGPR like *P. fluorescens* (NBAIL-PFDWD) is very limited. The study has shown that *P. fluorescens* expressed a wide array of stress response genes which ameliorate its survival in environmental perturbations. The stress response genes identified are complex and *P. fluorescens* does possess unique pathways to survive under osmotic stress. The culture is being promoted as a useful biological control agent in stressed soils. Application of stress tolerant strains of microorganisms could be considered as a feasible choice in stress prone areas (Meena *et al.*, 2017).

## ACKNOWLEDGMENT

The authors are grateful to The World Bank and Indian Council for Agricultural Research (ICAR) for funding the Research under the National Agricultural Innovation Project (NAIP).

## REFERENCES

- Ashwitha K, Rangeshwaran R, Vajid NV, Sivakumar G, Jalali SK, Rajalakshmi K and Manjunath H. 2013. Characterization of abiotic stress tolerant *Pseudomonas* spp. occurring in Indian soils. *J Biol Control*. **27**: 319–328.

- Atia A, Alrawaiq N, Abdullah A. 2014. A review of NAD(P) H:Quinone oxidoreductase 1 (NQO1); A multifunctional antioxidant enzyme. *J Appl Pharma Sci.* **4**: 118–122.
- Auriol A, Bestel-Corre G, Claude J, Soucaille P, Meynial-Salles I. 2011. Stress-induced evolution of *Escherichia coli* points to original concepts in respiratory cofactor selectivity. *PNAS* **108**: 1278–1283. <https://doi.org/10.1073/pnas.1010431108> PMID:21205901 PMCid:PMC3029715
- Benitez-Paez A, Villarayo M and Armengod. 2012. The *Escherichia coli* RlmN methyltransferase is a dual-specificity enzyme that modifies both rRNA and tRNA and controls translational accuracy. *RNA* **18**: 1783–1795. <https://doi.org/10.1261/rna.033266.112> PMID:22891362 PMCid:PMC3446703
- Beyenback KW, Wiezoerck H. 2006. The V-type H<sup>+</sup> ATPase: molecular structure and function, physiological roles and regulation. *J Exp Biol.* **209**: 577–589. <https://doi.org/10.1242/jeb.02014> PMID:16449553
- Boor KJ. 2006. Bacterial stress responses: What doesn't kill them can make them stronger. *PLoS Biol.* **4**: 0018–0023.
- Borovinskaya MA, Pai RD, Zhang W, Schuwirth BS, Holton JM, Hirokawa G, Kaji H, Kaji A. and Cate JHD. 2007. Structural basis for aminoglycoside inhibition of bacterial ribosome recycling. *Nat Struct Mol Biol.* **14**: 727–732. <https://doi.org/10.1038/nsmb1271> PMID:17660832
- Caldas TD, Yaagoubi AE, Richarme G. 1998. Chaperone properties of bacterial elongation factor EF-Tu. *J Biol Chem.* **273**: 11478–11482. <https://doi.org/10.1074/jbc.273.19.11478>. PMID:9565560
- Cameron A, Fridrich E, Huynh S, Parker CT and Gaynor EC. 2012. Hyperosmotic stress response of *Campylobacter jejuni*. *J Bacteriol.* **194**: 6116–6130. <https://doi.org/10.1128/JB.01409-12> PMID:22961853 PMCid:PMC3486418
- Chen M, Cao J, Zheng C, Liu Q. 2006. Directed evolution of an artificial bifunctional enzyme, c-glutamyl kinase/c-glutamyl phosphate reductase, for improved osmotic tolerance of *Escherichia coli* transformants. *FEMS Microbiol Lett.* **263**: 41–47. <https://doi.org/10.1111/j.15746968.2006.00397.x> PMID:16958849
- Cohen BE. 2014. Functional linkage between genes that regulate osmotic stress responses and multidrug resistance transporters: Challenges and opportunities for antibiotic discovery. *Antimicrob Agents Chemother.* **58**: 640–646. <https://doi.org/10.1128/AAC.02095-13> PMID:24295980 PMCid:PMC3910827
- Csonka LN. 1989. Physiological and genetic responses of bacteria to osmotic stress. *Microbiol Rev.* **53**: 121–147. PMID:2651863 PMCid:PMC372720
- Feng G, Tsui HT, Winkler ME. 1996. Depletion of the cellular amounts of the MutS and MutH methyl-directed mismatch repair proteins in stationary-phase *Escherichia coli* K-12 cells. *J Bacteriol.* **178**: 2388–2396. <https://doi.org/10.1128/jb.178.8.2388-2396.1996> PMID:8636043 PMCid:PMC177950
- Freeman BC, Chen C and Beattie GA. 2010. Identification of the trehalose biosynthetic loci of *Pseudomonas syringae* and their contribution to fitness in the phyllosphere. *Environ Microbiol.* **12**: 1486–1497. PMID:20192963
- Freeman BC, Chen C, Yu X, Nielsen X, Peterson, Beattie GA. 2013. Physiological and transcriptional responses to osmotic stress of two *Pseudomonas syringae* strains that differ in epiphytic fitness and osmotolerance. *J Bacteriol.* **195**: 4742–4752. <https://doi.org/10.1128/JB.00787-13> PMID:23955010 PMCid:PMC3807433
- Harris RS, Feng G, Ross KJ, Sidhu R, Thulin C, Longerich S, Sztigetly SK, Winkler ME, Rosenberg SM. 1997. Mismatch repair protein MutL becomes limiting during stationary-phase mutation. *Genes Dev.* **11**: 2426–2437. <https://doi.org/10.1101/gad.11.18.2426> PMID:9308969 PMCid:PMC316514
- Janoshi L, Shimizu I, Kaji A. 1994. Ribosome recycling factor (ribosome releasing factor) is essential for bacterial growth. *Proc Natl Acad Sci.* **91**: 4249–4253. <https://doi.org/10.1073/pnas.91.10.4249>
- Jensen KF, Larsen S. 2003. Dihydroorotate dehydrogenase of *Escherichia coli*. pp.11–21 In: Selinsky, B.S. (Eds.), *Membrane protein protocols; expression, purification and characterization*. Humana Press, Towata, NJ. <https://doi.org/10.1385/1-59259-400-X:11>
- Karas VO, Westerlaken I, Meyer AS. 2015. The DNA-binding protein from starved cells (Dps) utilizes dual functions to defend cells against multiple stresses. *J Bacteriol.* **197**: 3206–3215. <https://doi.org/10.1128/JB.00475-15> PMID:26216848 PMCid:PMC4560292
- Kawasaki S, Satoh T, Todoroki M, Niimura Y. 2009. b-Type dihydroorotate dehydrogenase is purified as a H<sub>2</sub>O<sub>2</sub>-forming NADH oxidase from *Bifidobacterium bifidum*. *Appl Environ Microbiol.* **75**: 629–636. <https://doi.org/10.1128/AEM.02111-08> PMID:19060157 PMCid:PMC2632149
- Kets EPW, Galinski EA, de Wit M, de Bont JAM, Heipieper HJ. 1996. Mannitol, a novel bacterial compatible sol-

- ute in *Pseudomonas putida* S12. *J Bacteriol.* **178**: 6665–6670. <https://doi.org/10.1128/jb.178.23.6665-6670>. PMID:8955280 PMCID:PMC178559
- Kuhlmann AU, Hoffmann T, Bursy J, Jebbar M, Bremer E. 2011. Ectoine and hydroxyectoine as protectants against osmotic and cold stress: Uptake through the SigB-controlled betaine-choline-carnitine transporter-type carrier EctT from *Virgibacillus pantothenicus* *J Bacteriol.* **193**: 4699–4708. <https://doi.org/10.1128/JB.05270-11> PMID:21764932 PMCID:PMC3165649
- Kusuya Y, Kurokawa K, Ishikawa S, Ogasawara N, Oshima T. 2011. Transcription factor GreA contributes to resolving promoter-proximal pausing of RNA polymerase in *Bacillus subtilis* cells *J Bacteriol.* **193**: 3090-3099. <https://doi.org/10.1128/JB.00086-11> PMID:21515770 PMCID:PMC3133182
- Larsen NA, Nash TJ, Morningstar M, Shapiro AB, Joubran C, Blackett CJ, Patten AD, Boriack- Sjodin A, Doig P. 2012. An aminoquinazoline inhibitor of the essential bacterial cell wall synthetic enzyme GlmU has a unique non-protein-kinase-like binding mode. *Biochem J.* **446**: 405–413. <https://doi.org/10.1042/BJ20120596> PMID:22721802
- Lau SKP, Fan RYY, Ho TCC, Wong GKM, Tsang AKL, Teng JLL, Chen W, Watt RM, Curreem SOT, Tse H, Yuen K, Woo PCY. 2011. Environmental adaptability and stress tolerance of *Laribacter hongkongensis*: a genome-wide analysis. *Cell Biosci.* **1**: 22–49. <https://doi.org/10.1186/2045-3701-1-22> PMID:21711489 PMCID:PMC3135505
- Lenhart JS, Brandes ER, Schroeder JW, Sorenson RJ, Showalter HD, Simmons LA. 2014. RecO and RecR are necessary for RecA loading in response to DNA damage and replication fork stress. *J Bacteriol.* **196**: 2851–2860. <https://doi.org/10.1128/JB.01494-14> PMID:24891441 PMCID:PMC4135682
- Li K, Jiang T, Yu B, Wang L, Gao C, Ma C, Xu P, Ma Y. 2012. Transcription elongation factor GreA has functional chaperone activity. *PLoS One* **7**: 47521-47527. <https://doi.org/10.1371/journal.pone.0047521> PMID:23251328 PMCID:PMC3521015
- Liang X, Zhang L, Natarajan SK, Becker DF. 2013. Proline mechanisms of stress survival. *Antioxid Redox Signal.* **19**: 998-1011. <https://doi.org/10.1089/ars.2012.5074> PMID:23581681 PMCID:PMC3763223
- Lindgren JK, Thomas VC, Olson ME, Chaudhari SS, Nuxoll AS, Schaeffer CR, Lindgren KE, Jones J, Zimmerman MC, Dunman PM, Bayles KW, Fey PD. 2014. Arginine deaminase in *Staphylococcus epidermidis* functions to augment biofilm maturation through pH homeostasis. *J Bacteriol.* **196**: 2277–2289. <https://doi.org/10.1128/JB.00051-14> PMID:24727224 PMCID:PMC4054176
- Liu X, Basu U, Miller P, McMullen L. 2014. Stress and adaptation of *Listeria monocytogenes* 08-5923 exposed to a sub-lethal dose of carboxycyclin A. *Appl Environ Microbiol.* **80**: 3835-3841. <https://doi.org/10.1128/AEM.00350-14> PMID:24747893 PMCID:PMC4054213
- Louis P, Galinski EA. 1997. Characterization of genes for the biosynthesis of the compatible solute ectoine from *Marinococcus halophilus* and osmoregulated expression in *Escherichia coli*. *Microbiol.* **143**: 1141-1149. <https://doi.org/10.1099/00221287-143-4-1141> PMID:9141677
- Lovett ST. 2011. The DNA exonucleases of *Escherichia coli*. *EcoSal. Plus.* doi:10.1128/ecosalplus.4.4.7. <https://doi.org/10.1128/ecosalplus.4.4.7>
- Lusetti SL, Cox MM. 2002. The bacterial RecA protein and the recombinational DNA repair of stalled replication forks. *Ann Rev Biochem.* **71**: 71-100. <https://doi.org/10.1146/annurev.biochem.71.083101.133940> PMID:12045091
- Macomber L, Elseby SP, Hausinger RP. 2011. Fructose-1,6-bisphosphate aldolase (class II) is the primary site of nickel toxicity in *Escherichia coli*. *Mol Microbiol.* **82**: 1291-1300. <https://doi.org/10.1111/j.1365-2958.2011.07891.x> PMID:22014167 PMCID:PMC3225726
- Maguire M, Coates ARM, Henderson B. 2002. Chaperonin 60 unfolds its secrets of cellular communication. *Cell Stress Chaperones* **7**: 317–329. <https://doi.org/10.1379/1466-1268%282002%29007%3C0317:CUISOC%3E2.0.CO%3B2>
- Mande SC, Kumar SCM. 2013. Evolution of bacterial chaperonin 60 paralogues and moonlighting activity. pp. 101-121. In: Henderson, B. (Ed.), *Moonlighting cell stress proteins in microbial infections, heat shock proteins*. Springer Science, Dordrecht. [https://doi.org/10.1007/978-94-007-6787-4\\_7](https://doi.org/10.1007/978-94-007-6787-4_7)
- Marquis RE, Bender GR, Murray DR, Wong A. 1987. Arginine deiminase system and bacterial adaptation to acid environments. *Appl Environ Microbiol.* **53**: 198–200. PMID:3103530 PMCID:PMC203628
- Meena KK, Sorty AM, Bitla UM, Choudhary K, Gupta P, Pareek A, Singh DP, Prabha R, Sahu PK, Gupta VK, Singh HB, Krishanani KK, Minhas PS. 2017. Abiotic stress responses and Microbe-mediated mitigation in plants: The omics strategies. *Fron Plant*

- Sci.* **8**:172. <https://doi.org/10.3389/fpls.2017.00172> PMID:28232845 PMCID:PMC5299014
- Mochalkin I, Lightle S, Zhu S, Ohren JF, Spessard C, Chirgadze NY, Banotai C, Melnick M, McDowell L. 2007. Characterization of substrate binding and catalysis in the potential antibacterial target N-acetylglucosamine-1-phosphate uridylyltransferase (GlmU). *Prot Sci.* **16**: 2657–2666. <https://doi.org/10.1110/ps.073135107> PMCID:PMC2222810
- Msadek T, Dartois V, Kunst F, Herbaud ML, Denizot F, Rapoport G. 1998. ClpP of *Bacillus subtilis* is required for competence development, motility, degradative enzyme synthesis, growth at high temperature and sporulation. *Mol Microbiol.* **27**: 899–914. <https://doi.org/10.1046/j.1365-2958.1998.00735.x> PMID:9535081
- Nurenberg E, Tampe R. 2013. Tying up loose ends: ribosome recycling in eukaryotes and archaea. *Trends Biochem Sci.* **38**: 64–74. <https://doi.org/10.1016/j.tibs.2012.11.003> PMID:23266104
- Oliver A, Baquero F, Blazquez J. 2002. The mismatch repair system (mutS, mutL and uvrD genes) in *Pseudomonas aeruginosa*: molecular characterization of naturally occurring mutants. *Mol Microbiol.* **43**:1641–1650. <https://doi.org/10.1046/j.1365-2958.2002.02855.x> PMID:11952911
- Pastor M, Salvador M, Argandoña M, Bernal V, Reina-Bueno M, Csonka LN, Iborra LJ, Vargas C, Nieto NJ, Cánovas M. 2010. Ectoines in cell stress protection: Uses and biotechnological production. *Biotech Adv.* **28**: 782–801. <https://doi.org/10.1016/j.biotechadv.2010.06.005> PMID:20600783
- Paul D, Dineshkumar N, Nair S. 2006. Proteomics of a plant growth promoting rhizobacterium, *Pseudomonas fluorescens* MSP-393, subjected to salt shock. *World J Microbiol Biotechnol.* **22**: 369–374. <https://doi.org/10.1007/s11274-005-9043-y>
- Perez-Arellano I, Carmona-Alvarez F, Martínez AI, Rodríguez-Díaz J, Cervera J. 2010. Pyrroline-5-carboxylate synthase and proline biosynthesis: From osmotolerance to rare metabolic disease. *Prot Sci.* **1**: 372–382. <https://doi.org/10.1002/pro.340>
- Rangeshwaran R, Ashwitha K, Sivakumar G, Jalali SK. 2013. Analysis of proteins expressed by an abiotic stress tolerant *Pseudomonas putida* (NBAIL-RPF9) isolate under saline and high temperature conditions. *Curr Microbiol.* **67**: 659–667. <https://doi.org/10.1007/s00284-013-0416-4> PMID:23828182
- Ryzhikov M, Koroleva O, Postnov D, Tran A, Korolev S. 2011. Mechanism of RecO recruitment to DNA by single-stranded DNA binding protein. *Nucleic Acids Res.* **1**: 1–10. <https://doi.org/10.1093/nar/gkr199>
- Sandhya V, Ali SkZ, Grover M, Reddy G, Venkateswarlu B. 2009. Alleviation of drought stress effects in sunflower seedlings by exopolysaccharides producing *Pseudomonas putida* strain P45. *Biol Fert Soil.* **46**: 17–26. <https://doi.org/10.1007/s00374-009-0401-z>
- Schelin J, Lindmark F, Clarke A.K. 2002. The clpP multigene family for the ATP-dependent Clp protease in the cyanobacterium *Synechococcus*. *Microbiol.* **148**: 2255–2265. <https://doi.org/10.1099/00221287-148-7-2255> PMID:12101312
- Song L, Wang G, Malhotra A, Deutscher MP, Liang W. 2016. Reversible acetylation on Lys501 regulates the activity of RNase II. *Nucleic Acids Res.* **44**: 1979–1988. <https://doi.org/10.1093/nar/gkw053> PMID:26847092 PMCID:PMC4797298
- Starosta AL, Lassak E, Jung K, Wilson DN. 2014. The bacterial translation stress response. *FEMS Microbiol Rev.* **38**: 172–1201. <https://doi.org/10.1111/1574-6976.12083> PMID:25135187 PMCID:PMC4227928
- Takahagi M, Iwasaki H, Nakata A, Shinagawa H. 1991. Molecular analysis of the *Escherichia coli* ruvC gene, which encodes a Holliday junction-specific endonuclease. *J Bacteriol.* **173**: 5747–5753. <https://doi.org/10.1128/jb.173.18.5747-5753.1991> PMID:1885548 PMCID:PMC208306
- Torrents E, Grinberg I, Gorovitz B, Lundström H, Borovok I, Aharonowitz Y, Sjöberg B, Cohen, G. 2007. NrdR controls differential expression of the *Escherichia coli* ribonucleotide reductase genes. *J Bacteriol.* **189**: 5012–5021. <https://doi.org/10.1128/JB.00440-07> PMID : 17496099 PMCID : PMC1951866
- Udaondo Z, Duque E, Fernández M, Molina L, Torre J, Bernal P, Niqui J, Pini C, Roca A, Matilla MA, Molina-Henares MA, Silva-Jimenez H, Navarro-Aviles G, Busch A, Lacal J, Krell T, Segura A and Ramos J. 2012. Analysis of solvent tolerance in *Pseudomonas putida* DOT-T1E based on its genome sequence and a collection of mutants. *FEBS Lett.* **586**: 2932–2938. <https://doi.org/10.1016/j.febslet.2012.07.031> PMID:22819823
- Villarroya M, Perez-Roger I, Macià F, Armengod ME. 1998. Stationary phase induction of dnaN and recF, two genes of *Escherichia coli* involved in DNA replication and

KARKERA ASHWITHA *et al.*

repair. *EMBO J.* **17**: 1829–1837. <https://doi.org/10.1093/emboj/17.6.1829> PMID:9501104 PMCID:PMC1170530

Wood JM. 1999. Osmosensing by bacteria: Signals and membrane-based sensors. *Microbiol Mol Biol Rev.* **63**: 230-262. PMID:10066837 PMCID:PMC98963

Wood JM. 2015. Bacterial responses to osmotic challenges. *J Gen Physiol.* **145**: 381-388. <https://doi.org/10.1085/jgp.201411296> PMID:25870209 PMCID:PMC4411257

org/10.1085/jgp.201411296%20PMid:25870209%20PMCID:PMC4411257

Zhang YM, Rock CO. 2008. Acyltransferases in bacterial glycerophospholipid synthesis. *J Lipid Res.* **49**: 1867–1874. <https://doi.org/10.1194/jlr.R800005-JLR200> PMID:18369234 PMCID:PMC2515527