



Relaxant and Antispasmodic Activities of Aqueous Extract from *Thymus algeriensis* Boiss. and Reut.

Leila Beyi¹, Hassan Zrouri¹, Hanane Makrane^{1,2}, Hassane Mekhfi¹, Abderrahim Ziyat¹, Mohamed Bnouham¹ and Mohammed Aziz^{1*}

¹Laboratory of Bioresources, Biotechnologies, Ethnopharmacologie et Santé, Faculty of Sciences, Mohamed the First University, PB 717, 60000, Oujda, Morocco; azizmo5@yahoo.fr

²Research Team: Biology, Environment and Sustainable Development, Department of Matter and Life Sciences, Ecole Normale Supérieure, Abdelmalek Essaadi University, Martil-Tetouan, Morocco

Abstract

Thymus algeriensis have been vastly utilized for intestinal disorders. The purpose of this investigation was to scrutinize the probable mechanism for its utilization in the spasm disorder. Aqueous extract of this medicinal plant (AqTA) was tested *in vitro* on rat and rabbit jejunum. The extract produced relaxation of rabbit jejunum. This relaxation does not depend on the adrenergic pathway, the AqTA induces inhibition irrespective of the presence or absence of adrenergic inhibitors. AqTA engendered a concentration-dependent (0.1-5 mg/ml) relaxation of carbamylcholine chloride (CCh) and K⁺ provoked tones in rat intestine with IC₅₀ values of 2.06 ± 0.26 and 3.55 ± 0.48 (mg/ml) respectively. This extract likewise induced a dose-dependent (0.1-3 mg/ml) rightward shift in the CCh and Ca⁺⁺ dose-response curves. The AqTA alone has decreased more significantly the percentage of contraction of rat jejunum than the AqTA pre-incubated with atropine or hexamethonium then contracted with KCl; but there is no significant difference by those pre-incubated with methylene blue or L-NAME. When the intestine was pretreated with nifedipine and contracted by CCh, the antispasmodic effect provoked by AqTA with and without pre-incubation with nifedipine is statistically not significant. In conclusion AqTA acts possibly on the voltage dependent Ca⁺⁺ channel and cholinergic receptors but did not act on adrenergic receptors, NO and guanylatecyclase pathway. This investigation may explicate some of its traditional utilization in gut illnesses.

Keywords: Antispasmodic, Jejunum, Relaxant, Smooth muscle, *Thymus algeriensis*

1. Introduction

Thymus algeriensis Boiss. and Reut. is an endemic species found in African countries along the Mediterranean Sea. It belongs to the *Thymus* genus. Numerous *Thymus* species are greatly employed in culinary and medicine. In Morocco *Thymus algeriensis*, known locally under the common name “Zaatar”, is found in the Atlas and Rif mountains and in the BéniSnassen Forest. It is a perennial plant with 4 to 7 mm long internodes emerging as a tuft from the short, woody stump¹. Its active principles are gallic acid, catechin, epicatechin, coumaric acid, apigenin, naringenin, rutin, quercetin, kaempferol, tannins, rutin², flavonol, flavone glycoside derivatives and phenolic acids,

specially rosmarinic acid and kaempferol-O-glucuronide³. The volatile components varied among populations; at least five chemotypes according to main compounds have been found: -pinene/ 1,8-cineole / -pinene, 1,8-cineole/ caryophyllene oxide/ camphor, linalool and thymol chemotypes⁴. In North Africa traditionally it is used to cure the problems of respiration, digestion and against abortion^{5,6}. Previous studies have shown that it had antioxidant, antimicrobial, healing gastric ulcers^{7,8} and antitumor⁹ activities.

In a preceding work, we obtained an antidiarrheal action of the aqueous extract of the *T. algeriensis*¹⁰. The purpose of this report is to continue this work on the digestive tract problems and to attempt to elucidate how

*Author for correspondence

this extract intervenes on the relaxation pathways of the muscle fibers of rabbit and rat jejunums.

2. Materials and Methods

2.1 Solutions and Drugs

For twenty years we have always used the same solutions for antispasmodic tests. See Normal Krebs-Henseleit Buffer (KHB), Calcium-free high K⁺, High K⁺ KHB (75 mM) and Calcium-free KHB solutions composed of (in mM) in the reference¹¹. The drugs employed in spasmolytic tests were L-NAME from Calbiochem, methylene blue from Sigma-Aldrich, carbamylcholine chloride (CCh), prazosin, nifedipine, yohimbine, propranolol were purchased from Sigma chemical co, hexamethonium from Across organics, atropine from Fluka.

2.2 HPLC Analysis of Phenolic Compounds

HPLC analysis was performed to analyze the phenolic compounds present in the plant¹².

2.3 Plant Material

T.algeriensis was harvested from Oujda province in the spring of 2014, was authenticated by Prof. Benyounes Haloui and a voucher specimen was deposited in the Herbarium of the Sciences Faculty, University Mohammed the First, Oujda, Morocco under the reference number (HUMPOM 425).

2.4 Preparation of the Extract

Inspired by the Moroccan tradition, 50g of the aerial part of *T. algeriensis* (AqTA) was boiled in 1l of water for 15

min and dehydrated to dryness to give a crude residue (yields: 18%).

2.5 Animals

Female and male New Zealand rabbits (1-2 Kg) and Wistar rats (200-300kg) were provided from the animal house of the Sciences Faculty in Oujda (Morocco). Animals had free access to water but food was retiring 18h prior to experimentation. All animals were treated for in conformity with the internationally accepted Guide for the care and use of laboratory animals, published by US National Institutes of Health (NIH Publication N°. 85-23, revised in 1985).

2.6 Spasmolytic Study

The spasmolytic activity of the jejunums was tested exactly like the one we studied on *Origanum majorana*¹¹.

2.7 Statistics

The results are expressed as means ± S.E.M. The statistical significance of data was analyzed using Student's t-test, P<0.05 was considered as significant. The 50% inhibitory concentration (IC₅₀) was determined by linear regression method.

3. Results

3.1 HPLC

HPLC analysis of the phenolic composites showed that the AqTA contains syringic acid, rutin, coumaric acid, luteolin, cinnamic acid, apigenin, and quercetin (Table 1, Figure 1).




Figure 1. HPLC chromatogram of aqueous extract from *Thymus algeriensis*.

3.2 Effects of AqTA on Rabbit Intestine

Contact of the intestines from 0 to 5 mg/ml of the AqTA decreased the mean magnitude of the spontaneous tonus to almost total inhibition of control with an IC_{50} value of 1.88 ± 0.07 mg/ml (Figure 2). This action was reversible as the intestine recovered its basal activity after rinsing twice with KHB. Adrenaline at 10^{-6} M inhibited rabbit jejunum contractions. The addition to the KHB medium of β (Propranolol), α_2 (Yohimbine) and α_1 (Prazosin) adrenergic inhibitors with AqTA provoked inhibition of the spontaneous contractions of the rabbit jejunum.

Table 1. Results of HPLC analysis Aqueous extract from *Thymus algeriensis* compared to Retention time of 18 standard phenol acids and flavonoids

Standards	RT(min)	AqTA
Chlorogenic acid	9.23	-
Catechin	9.33	-
3,4-dihydropyran	9.60	-
4-hydro phenyl	10.27	-
Vanillic acid	10.50	-
4-hydroxybenzoic acid	10.62	-
Syringic acid	10.76	+
Caffeic acid	11.11	-
Vanillin	11.35	-
p-coumaric acid	12.73	-
trans-ferulic acid	13.35	-
Rutin	15.18	+
Coumaric acid	15.47	+
Luteolin	16.14	+
Cinnamic acid	16.72	+
Apigenin	17.46	+
Quercetin	17.72	+
trans-chalcone	20.65	-

“+” sign indicates the presence of phenolic compounds;

“-” sign indicates the absence of phenolic compounds.

3.3 Inhibition of Dose-response to Carbachol and $CaCl_2$

AqTA caused a dose-dependent (0.1-5mg/ml) inhibition CCh and K^+ induced tones of rat jejunum preparations with IC_{50} values of 2.06 ± 0.26 and 3.55 ± 0.48 (mg/ml) respectively (Figures 3 and 4). Our extract likewise caused a dose-dependent (0.1-3 mg/ml) rightward shift in the CCh (Figure 5) and Ca^{++} dose-response curves (Figure 6).




Figure 2. Effect of AqTA on rabbit jejunum preparations. Values shown are mean \pm S.E.M. of six experiments (***) $P < 0.001$.




Figure 3. Effect of aqueous extract from *Thymus algeriensis* on Carbachol (10^{-6} M) induced contractions of rat jejunum preparation. Values shown are mean \pm S.E.M. of six experiments. (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$).




Figure 4. Effect of aqueous extract from *Thymus algeriensis* on KCl (75mM) induced contractions of rat jejunums. Values shown are mean \pm S.E.M. of six experiments. (** $P < 0.01$; *** $P < 0.001$).




Figure 5. Concentration-response curves of carbachol (CCh) in the presence of different concentrations of AqTA (0; 0.3; and 3 (mg/ml)). Values shown are mean \pm S.E.M. of six experiments. (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$).




Figure 6. Concentration-response curves of CaCl_2 in the presence of different concentrations of AqTA (0; 0.3; 1 and 3 (mg/ml)). Values shown are mean \pm S.E.M. of six experiments. (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$).

3.4 Effect of AqTA with Different Pharmacological Blocking Agents on Rat Jejunum Pre-contracted by KCl or CCh

We compared the action of 3 mg/ml of AqTA, concentration approaching the IC_{50} , with pharmacological blocking agents habitually utilised to reduce the contraction provoked by KCl. The AqTA alone has decreased more significantly the percentage of contraction of rat jejunum than the AqTA pre-incubated with atropine or hexamethonium; but there is no significant difference

by those pre-incubated with L-NAME or methylene blue (Figure 7A).

When KCl was replaced by Carbachol, the antispasmodic effect provoked by AqTA with and without pre-incubation with nifedipine is statistically not significant (Figure 7B).




Figure 7. Effect of aqueous extract of *Thymus algeriensis* (3 mg/ml) on contractions of the rat jejunum preincubated with hexamethonium (10^{-4} M), atropine (10^{-6} M), methylene blue (10^{-5} M) and L-NAME (10^{-4} M) for 20 min, and then pre-contracted by 75 mM KCl. (A) or pre-incubated with nifedipine (10^{-6} M) for 20 min, and then pre-contracted by CCh 10^{-6} M (B). Values shown are mean \pm S.E.M. of six experiments. (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$). NS = Not significant. (***) compared to the control and # and NS compared to the extract without inhibitors).

4. Discussion

Due to the traditional reputation as antispasmodic, aqueous extract of *Thymus algeriensis* was tested for its possible spasmolytic effect in spontaneously contracting rabbit jejunum preparations because it's easier to quantify relaxation than in spontaneous contraction of rat jejunum. It is known that intestinal smooth muscle contracts rhythmically in the absence of hormonal and neuronal stimulations; such contractions are referred to as phasic¹³ and are commenced by the act of the interstitial cell of Cajal¹⁴. AqTA inhibited spontaneous contractions in a reversible dose dependent manner. Sympathetic nerves of the autonomic nervous system regulate intestine motility in a round-about way across the enteric nervous system, making an inhibitory action on motility¹⁵. We wanted to know if our extract acts on the adrenergic pathway. For this, we blocked adrenergic receptors, β by the propranolol, α_2 by the yohimbine and α_1 by the prazosin (put all together). We remarked that noradrenaline does not have any impact on the tensions; whereas the extract at 5mg/ml, the dose formerly described to provide a manifest relaxant effect, generates the prevention of these contractions. Like extracts from other plants that we have tested in our laboratory, as *Cistus ladaniferus*¹⁶ and *Origanum majorana*¹¹, AqTA also does not act via adrenergic receptors.

Usually antispasmodic action of medicinal plants is due to Ca^{++} channel blockade^{17,18}. To check if the antispasmodic action of the AqTA is as well due via equivalent process, this extract was evaluated on high K^+ -induced contraction. This latter provokes a depolarization of the plasma membrane and subsequently the voltage-gated Ca^{++} channels (VGCaCs) open and engendered Ca^{++} get in hyaloplasm¹⁹. A chemical which can prevent K^+ provoked tone is hence reputed to be a calcium channel blocking²⁰. Therefore, antispasmodic effect of rich K^+ medium (75 mM) –induced tone of rat intestine by AqTA may reflect the restrained Ca^{++} entrance through voltage-dependent channels. This supposition was in addition fortified during the pre-treatment of the gut with *T. algeriensis* which produced a concentration-dependent rightward shift in the concentration-response of $CaCl_2$ ²¹. AqTA apparently does not act on L-type VGCaCs, it could intervene in other types of VGCaCs.

The cholinergic receptor pathway was tested using Carbachol which is a structural analogue of acetylcholine^{22,23}. AqTA inhibited by a dose-dependent mode, the tone generated with this analogue. This purpose was reinforced when pre-treatment of the jejunum with thym extract engendered a concentration-dependent

rightward shift in the concentration-response of CCh. To know which cholinergic receptors may be involved, we used the muscarinic receptor antagonist atropine^{24,25} and the nicotinic receptor antagonist hexamethonium^{26,27}. Atropine, but not hexamethonium, have impaired the initial relaxed aftermath of AqTA. Subsequently it can be proposed that our plant intervenes directly on the muscarinic receptors.

Nitric oxide (NO), a transmitter gut nerves in the myenteric plexus nerves in the myenteric plexus throughout the gastrointestinal tract²⁸, was able to relax intestine²⁹. It relaxes smooth muscle fibers via activation of dissolvable guanylate cyclase^{30,31} and an increase in intracellular cGMP. We wanted to know if AqTA acts via the NO and/or that of guanylatecyclase. The employment of L-NAME, an inhibitor of NO and blue methylene, a blocker agent of guanylatecyclase pathways³², showed that AqTA did not act on these two pathways.

The relaxant and antispasmodic effect can be due to Apeginin³³, Luteolin^{34–36} and Quercetin^{37–39}, known by their antispasmodic effect and that are existing in AqTA. These components could act alone or together with other unidentified compounds.

5. Conclusion

These results suggest that the spasmolytic activity of AqTA are mediated possibly through Ca^{++} antagonist mechanism and cholinergic receptors but did not act on adrenergic receptors, NO and guanylate cyclase. This study may explain some of the traditional uses of the plant in gastrointestinal disorders, and with our previous work¹⁰ reinforce positively the practice of Thym in Moroccan folk medicine as a spasmolyticgut agent.

6. Acknowledgement

Our thanks to Badraoui M and Ramdaoui K for supplying us with experimental animals and the Sciences faculty of Oujda for supporting this research.

7. References

1. Benabid A. Flore et écosystèmes du Maroc. Évaluation et préservation de la biodiversité. Paris : Édition Ibis Press; 2000. p. 159–61.
2. Guesmia F, Saidib I, Bouzennab H, Hfaiedhb N, Landoulsi A. Phytocompound variability, antioxidant and antibacterial activities, anatomical features of glandular and aglandular hairs of *Thymus hirtus* Willd. Ssp. *algeriensis* Boiss. and Reut. over developmental stages. South African Journal of

- Botany. 2019; 127:234–43. <https://doi.org/10.1016/j.sajb.2019.09.004>
3. Ziani BEC, Heleno SA, Bachari K, Dias MI, Alves MJ, Barros L, Ferreira I CFR. Phenolic compounds characterization by LC-DAD-ESI/MSn and bioactive properties of *Thymus algeriensis* Boiss. and Reut. and *Ephedra alata* Decne. *Food Res Int.* 2019; 116:312–19. <https://doi.org/10.1016/j.foodres.2018.08.041>. PMID: 30716951
 4. Ali IBE, Guetat A, Boussaid M. Chemical and genetic variability of *Thymus algeriensis* Boiss. and Reut. (Lamiaceae), a North African endemic species. *Ind Crop Prod.* 2012; 40(1):277–84. <https://doi.org/10.1016/j.indcrop.2012.03.021>
 5. Bellakhdar J. La pharmacopée marocaine traditionnelle. Médecine arabe ancienne et savoirs populaires. Ibisspress; 1997. p. 349.
 6. Giweli AA, Džamić AM, Soković MD, Ristić MS, Marin PD. Chemical composition, antioxidant and antimicrobial activities of essential oil of *Thymus algeriensis* wild-growing in Libya. *Cent Eur J Biol.* 2013; 8(5):504–11. <https://doi.org/10.2478/s11535-013-0150-0>
 7. Guesmi F, Ben FM, Mejri M, Landoulsi A. In-vitro assessment of antioxidant and antimicrobial activities of methanol extracts and essential oil of *Thymus hirtus* sp. algeriensis. *Lipids Health Dis.* 2014; 13:114–25. <https://doi.org/10.1186/1476-511X-13-138>. PMID:25159191 PMCid:PMC4176582
 8. Guesmi F, Ali MB, Barkaoui T, Tahri W, Mejri M, Ben-Attia M, *et al.* Effects of *Thymus hirtus* sp. Algeriensis Boiss. & Reut. (Lamiaceae) essential oil on healing gastric ulcers according to sex. *Lipids Health Dis.* 2014; 13:138–50. <https://doi.org/10.1186/1476-511X-13-138>. PMID:25159191. PMCid:PMC4176582
 9. Nikolić M, Glamočlija J, Ferreira ICFR, Calhelha RC, Fernandes, Marković T, Marković D, Giweli A, Soković M. Chemical composition, antimicrobial, antioxidant and antitumor activity of *Thymus serpyllum* L., *Thymus algeriensis* Boiss. and Reut and *Thymus vulgaris* L. essential oils. *Ind Crops Prod.* 2014 Jan; 52:183–90. <https://doi.org/10.1016/j.indcrop.2013.10.006>
 10. Beyi L, Aziz M, Makrane H, Karim A, Bnouham M, Ziyayat A. Antidiarrhoeal activity of *Thymus algeriensis* boiss. & reut. aqueous extract on rats and mice. *Int J Pharm Pharm Sci.* 2015; 7(3):292–4.
 11. Makrane H, Aziz M, Mekhfi H, Ziyayat A, Bnouham M, Legssyer A, Gressier B, Eto B. Antispasmodic and Myorelaxant Activity of Organic Fractions from *Origanum majorana* L. on Intestinal Smooth Muscle of Rodents. *Eur J Med Plant.* 2018; 23(2):1–11. <https://doi.org/10.9734/EJMP/2018/41075>
 12. Makrane H, El messaoudi M, Melhaoui A, El mzibri M, Benbacer L, Aziz M. 2018a. Cytotoxicity of the aqueous extract and organic fractions from *Origanum majorana* on human breast cell lines MDA-MB-231 and human colon cell lines HT-29. *Adv Pharmacol Sci.* <https://doi.org/10.1155/2018/3297193>. PMID:30210537. PMCid: PMC6126111
 13. Weisbrodt NW. Regulation: Nerves and smooth muscle. In: *Gastrointestinal Physiology*, Johnson LR (ed.). Philadelphia, PA: Mosby Elsevier; 2007. p. 13–21. <https://doi.org/10.1016/B978-0-323-03391-6.50006-6>
 14. McHale N, Hollywood M, Sergeant G, Thornbury K. Origin of spontaneous rhythmicity in smooth muscle. *J Physiol.* 2006; 570:23–8. <https://doi.org/10.1113/jphysiol.2005.098376>. PMID:16239271. PMCid:PMC1464298
 15. Hansen MB. The enteric nervous system II: Gastrointestinal functions. *Pharmacol Toxicol.* 2003; 92:249–57. <https://doi.org/10.1034/j.1600-0773.2003.920601.x>. PMID:12787256
 16. Aziz M, Tab N, Karim A, Mekhfi H, Bnouham M, Ziyayat A, *et al.* Relaxant effect of aqueous extract of *Cistus ladaniferus* on rodent intestinal contractions. *Fitoterapia.* 2006; 77:425–8. <https://doi.org/10.1016/j.fitote.2006.05.015>. PMID:16815641
 17. Gilani AH, Khan A, Ghayur MN, Ali SF and Herzig J. Antispasmodic Effects of Rooibos Tea (*Aspalathu slinearis*) is Mediated Predominantly through K⁺-channel Activation. *Basic Clin Pharmacol Toxicol.* 2006;99:365–73. https://doi.org/10.1111/j.1742-7843.2006.pto_507.x. PMID:17076689
 18. Makrane H, Aziz M, Berrabah M, Mekhfi H, Ziyayat A, Bnouham M, *et al.* Myorelaxant Activity of essential oil from *Origanum majorana* L. on rat and rabbit. *J Ethnopharmacol.* 2019; 228:40–9. <https://doi.org/10.1016/j.jep.2018.08.036>. PMID:30205180
 19. Bolton TB. Mechanism of action of transmitters and other substances on smooth muscles. *Physiol Rev.* 1979; 59:606–718. <https://doi.org/10.1152/physrev.1979.59.3.606>. PMID:37533
 20. Godfraind T, Miller R, Wibo M. Calcium antagonism and calcium entry blockade. *Pharmacol Rev.* 1986:321–416.

21. Rojas A, Bah M, Rojas JI, Serrano V, Pacheco S. Spasmolytic activity of some plants used by the Otomi Indians of Querétaro (México) for the treatment of gastrointestinal disorders. *Phytomedicine*. 1999; 6:367–71. [https://doi.org/10.1016/S0944-7113\(99\)80061-0](https://doi.org/10.1016/S0944-7113(99)80061-0)
22. Hardman JG, Umbird LE, Gliman AG, Goodman G. The pharmacological basis of therapeutics. 10th ed McGraw Hill, New York; 2001.
23. Guata YS, Sarr A, MoctarDièye A, Faye B. Myorelaxant and antispasmodic effects of the aqueous extract of *Mitragyna inermis* barks on Wistar rat ileum. *Fitoterapia*. 2004;447–50. <https://doi.org/10.1016/j.fitote.2004.03.006>. PMID:15261381
24. Arunlakshana O, Schild HO. Some quantitative uses of drug antagonists. *Br J Pharmacol Chemotherapy*. 1959; 14:48–58. <https://doi.org/10.1111/j.1476-5381.1959.tb00928.x>. PMID:13651579. PMCid:PMC1481829
25. Gilani AH, Shaheen F, Christopoulos A, Mitchelson F. Interaction of ebeinone, an alkaloid from *Fritillaria imperialis*, at two muscarinic acetylcholine receptor subtypes. *Life Sci*. 1997; 60(8):535–44. [https://doi.org/10.1016/S0024-3205\(96\)00691-1](https://doi.org/10.1016/S0024-3205(96)00691-1)
26. Barlow RB, Franks M, Pearson JDM. A comparison of the affinities of antagonists for acetylcholine receptors in the ileum, bronchial muscle and iris of the guinea-pig. *British Journal of Pharmacology*. 1972;300–8. <https://doi.org/10.1111/j.1476-5381.1972.tb06875.x>. PMID:4405611. PMCid:PMC1666339
27. Spencer NJ, Walsh M, Smith TK. Purinergic and cholinergic neuroneuronal transmission underlying reflexes activated by mucosal stimulation in the isolated guinea pig ileum. *J Physiol*. 2000;321–31. <https://doi.org/10.1111/j.1469-7793.2000.t01-1-00321.x>. PMID:10639107. PMCid:PMC2269751
28. Bult H, Boeckxstaens GE, Pelckmans PA, Jordaens FH, Van Maercke YM, Herman AG. Nitric oxide as an inhibitory non-adrenergic non-cholinergic neurotransmitter. *Nature*. 1990; 24, 345(6273):346–7. <https://doi.org/10.1038/345346a0>. PMID:1971425
29. Gustafsson BI, Delbro DS. Tonic inhibition of small intestinal motility by nitric oxide. *J Auton Nerv Syst*. 1993; 44:179–87. [https://doi.org/10.1016/0165-1838\(93\)90030-X](https://doi.org/10.1016/0165-1838(93)90030-X)
30. Furchgott RF, Zawadzki JV. The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature*. 1980; 288:373–6. <https://doi.org/10.1038/288373a0>. PMID:6253831
31. Wollin MS, Wood KS, Igharro LJ. Guanylate cyclase from bovine lung. A kinetic analysis of the regulation of the purified soluble enzyme by protoporphyrin IX, heme and nitrosylheme. *J Biol Chem*. 1982; 257:13312–20. [https://doi.org/10.1016/S0021-9258\(18\)33448-3](https://doi.org/10.1016/S0021-9258(18)33448-3)
32. Komaitis ME. Composition of the essential oil of marjoram (*Origanum majorana* L.). *Food Chem*. 1992; 45:117–18. [https://doi.org/10.1016/0308-8146\(92\)90020-3](https://doi.org/10.1016/0308-8146(92)90020-3)
33. Lemmens-Gruber R, Marchart E, Rawnduzi P, Engel N, Benedek B, Kopp B. Investigation of the spasmolytic activity of the flavonoid fraction of *Achillea millefolium* L. on isolated guinea-pig ilea. *Arzneimittel-Forschung/Drug Research*. 2006; 56(8):582–6. <https://doi.org/10.1055/s-0031-1296755>. PMID:17009839
34. Gorzalczyk S, Moscatelli V, Ferraro G. Artemisia copa aqueous extract as vasorelaxant and hypotensive agent. *J Ethnopharmacol*. 2013; 148(1):56–61. <https://doi.org/10.1016/j.jep.2013.03.061>. PMID:23588093
35. Fleer H, Verspohl J. Antispasmodic activity of an extract from *Plantago lanceolata* L. and some isolated compounds. *Phytomedicine*. 2007; 14(6):409–15. <https://doi.org/10.1016/j.phymed.2006.05.006>. PMID:17298877
36. Engelbertz J, Lechtenberg M, Studt L, Hensel A, Verspohl EJ. Bioassay-guided fractionation of a thymol-deprived hydrophilic thyme extract and its antispasmodic effect. *J Ethnopharmacol*. 2012; 141(3):848–53. <https://doi.org/10.1016/j.jep.2012.03.025>. PMID:22465593
37. Lozoya X, Meckes M, Abou-Zaid M, Tortoriello J, Nozzolillo C, Arnason JT. Quercetin glycosides in *Psidium guajava* L. leaves and determination of a spasmolytic principle. *Arch Med Res*. 1994; 25(1):11–15.
38. Melzig MF, Pertz HH, Krenn L. Anti-inflammatory and spasmolytic activity of extracts from *Drosera Herba*. *Phytomedicine*. 2001; 8(3):225–9. <https://doi.org/10.1078/0944-7113-00031>. PMID:11417917
39. Krenn L, Beyer G, Pertz HH, Karall E, Kremser M, Galambosi B, et al. In vitro antispasmodic and anti-inflammatory effects of *Drosera rotundifolia*. *Arzneimittel-Forschung/Drug Research*. 2004; 54(7):402–5. <https://doi.org/10.1055/s-0031-1296991>. PMID:15344845