

Study of the Risk Factors for Coronary Artery Disease and their Correlation with Thyroid Hormone Profile amongst Women with ST Segment Elevation in Acute Myocardial Infarction

Neelima Chafekar¹, Mayur Babras² and Madhuri Kirloskar^{3*}

¹Professor, Department of General Medicine, Dr. Vasantrao Pawar Medical College, Hospital and Research Centre, Nashik – 422003, Maharashtra, India

²Former PG Resident, Department of General Medicine, Dr. Vasantrao Pawar Medical College, Hospital and Research Centre, Nashik – 422003, Maharashtra, India

³Associate Professor, Department of General Medicine, Dr. Vasantrao Pawar Medical College, Hospital and Research Centre, Nashik – 422003, Maharashtra, India; mskirloskar@gmail.com

Abstract

Background: Thyroid hormone has a major role in the cardiovascular function and cardiac hemodynamics as well as to maintain the cardiovascular homeostasis. The present study was planned to assess risk factors for coronary artery disease and their correlation with thyroid hormone profile amongst women with ST segment elevation in acute myocardial infarction. **Material and Methods:** A total number of 78 females having ST elevation myocardial infarction diagnosed through detailed clinical history and ECG evaluations were included after they satisfy the eligibility criteria. Patients were subsequently evaluated for presence of risk factors of ischemic heart disease such as age, marital status, parity, and menopause, use of oral contraceptives, family history, obesity, diabetes mellitus, hypertension, hypercholesterolemia and hypertriglyceridemia. Their thyroid hormone profiles (T3, T4, TSH) were done and were correlated with risk factors for ischemic heart disease. **Results:** A total 78 patient included in the study most of the study population belongs to the age group of 40 to 50 years (43.6 %). Hypertension was the most common clinical features amongst study population (59%) followed by Diabetes (50%). Most of the study population had Normal Thyroid status (57.69%) followed by Hypothyroidism (25.64%) and Hyperthyroidism (16.67%) and SERUM TSH is the most sensitive test for evaluation. Diabetes Mellitus was statistically significant risk factor (p value<0.05) in patients with STEMI with different levels of thyroid. **Conclusion:** Patients with coronary artery disease especially in the presence of other risk factors should be screened for thyroid dysfunction specially diabetes. We recommend tests for thyroid disorders in acute coronary syndrome can give predictor for risk of morbidity and mortality in those subjects.

Keywords: MI-Myocardial Infarction, Hyperthyroidism, Hypothyroidism, STEMI-ST- Segment Elevation Myocardial Infarction, TSH- Thyroid Stimulating Hormone

1. Introduction

The term “myocardial infarction” focuses on the myocardium (the heart muscle) and the changes that occur in it due to the sudden deprivation of circulating blood. The main change is necrosis (death) of myocardial tissue. The word “infarction” comes from the Latin “infarcire”

meaning “to plug up or cram”. It refers to the clogging of the artery. The criteria’s of Myocardial Infarction are based on symptoms, cardiac enzymes, ST-T changes on ECG, regional wall abnormality present on imaging and coronary angiography.

Ischemic Heart Disease (IHD) is the biggest killer of women globally. IHD causes 8.6 million deaths

*Author for correspondence

among women annually, a third of all deaths in women worldwide. Every year more women than men die due to Ischemic heart disease. Women in low and middle income countries are in a worse situation than men, experiencing higher proportion of IHD deaths than men¹. Women continue to be under represented in research on heart disease. Most of the studies conducted on IHD are based on male population and whether we should implement the same guidelines on women counterparts is an unanswered question². With more data from Women's Ischemia Syndrome Evaluation Study (WISE), as well as other new studies during the past several years, an evolving knowledge regarding sex differences in IHD has emerged. Women and men with heart disease tend to differ in their presenting symptoms, access to investigations, treatment and overall prognosis³.

Women present with more atypical symptoms than men like back pain, shortness of breath, burning in the chest, nausea, or fatigue, which makes the diagnosis more difficult. Risk factors for IHD vary between males and females⁴. Diabetes mellitus is a stronger IHD risk factor in women than in men. Hypertension is associated with a two to threefold increased risk for IHD in women. In women, low levels of high density lipoprotein are strong predictors of higher IHD risk than high levels of low density lipoprotein⁵. Studies have shown complex relationship between IHD risk, estrogen, menopause and serum cholesterol in women^{6,7}. Anti-estrogenic effect of tobacco and smoking increases the risk of IHD in premenopausal women⁸.

Studies have shown, in women cardiovascular risk profiles improve with increasing levels of physical activity⁹. In women central obesity is observed as one of the major risk factor for IHD¹⁰. Despite differences between the sexes in risk factors, presentation, and response to treatment, women in our country continue to receive similar treatments to men on the basis of trials that include mainly male participants.

Thyroid hormone has a major role in the cardiovascular function and cardiac hemodynamics well as to maintain the cardiovascular homeostasis¹¹⁻¹³. A slight change in thyroid status affects ventricular function, serum cholesterol levels, heart rate and rhythm, which increases risk of coronary artery disease and cardiovascular mortality¹⁴. Nevertheless, the relation between anomalous thyroid function and cardiovascular effects remains indistinct¹⁵.

Clinical features characteristic of hyperthyroidism like tachycardia, high cardiac output, increased myocardial contractility, systolic blood pressure, and basal Ometabolism, as well as tremors suggests a hyperadrenergic state. This is all due to the sensitivity to catecholamine compounds¹⁶⁻¹⁸. Hypothyroidism instead, seems to evoke a hypoadrenergic state due to the presence of bradycardia, reduced basal metabolism and cardiac output, and the intracellular catecholamine production from circulation, which has been found to be lower during hypothyroidism¹⁹⁻²¹.

This study was conducted to evaluate important risk factors contributing to the IHD in women and its association with thyroid hormone profile as it is one of the contributory factor which increases risk of IHD in women with underlying other risk factors, hence needs to be evaluated.

2. Aims and Objectives

- To study clinical profile and cardiovascular risk factors in females admitted with ST Elevation Myocardial Infarction (STEMI) in the Coronary Care unit.
- To evaluate thyroid hormone profile (T3, T4, TSH) in females with ST Elevation Myocardial Infarction (STEMI)
- To find if there is any correlation of thyroid hormone dysfunction with other risk factors in females with ST Elevation Myocardial Infarction (STEMI).

3. Materials and Methods

Study Design- Cross-sectional study

Study Setting- Intensive Care Unit of Dr. Vasantrao Pawar Medical College, Hospital and Research Centre, Nashik, Maharashtra, India

Duration of Study- August 2016 To September 2018

Study Population- 78 Female patients with ST Elevation Myocardial Infarction (STEMI) admitted to ICU in our Hospital were included in the study.

3.1 Sample Size Calculation Formula

Formula: $Z^2 \frac{p \cdot q}{L^2}$

L^2

Where Z= 1.96 (critical value)

p=proportion of the diseased p= 0.02

q= 1-p

L= margin of error = 5%

3.2 Eligibility Criteria

3.2.1 Inclusion Criteria

- Female patients, irrespective of age, admitted with new onset STEMI were included in the study.

(Presence of STEMI will be assessed by analysis of standard 12 lead ECG)

ECG criteria for st segment elevation myocardial infarction²²

- New ST elevation at the J point in two contiguous leads with the following cut points
 - > 0.1 mv in all leads (except V2-V3)
 - In leads v2 -v3 the following cut points apply
- >0.15 mv in women

3.2.2 Exclusion criteria

- Females with previous history of myocardial infarction
- Females with history of neck irradiation, history of thyroidectomy
- Females with chronic renal failure or other longstanding systemic illness
- Females on medications like amiodarone, iodides, lithium, and lugols iodine, antidopaminergic drugs
- Patients with valvular heart diseases
- Patients with congenital heart disease
- Patients with pericarditis / pericardial effusion
- Patients with aortic aneurysms
- Patients who have not given written informed consent

3.3 Methodology

A total number of 78 females having ST elevation myocardial infarction diagnosed through detailed clinical history and ECG evaluation were included after they satisfy the eligibility criteria. Only those patients giving valid informed consent were included in the study.

Female patients with ST Elevation Myocardial Infarction (STEMI) admitted to ICU in a Tertiary Care Hospital were included in the study. Presence of STEMI was assessed by analysis of standard 12 lead ECG. Patients were subsequently evaluated for presence of risk factors of ischemic heart disease such age, marital

status, parity, and menopause, use of oral contraceptives, family history, obesity, diabetes mellitus, hypertension, hypercholesterolemia and hypertriglyceridemia. Their thyroid hormone profile (T3, T4, TSH) was done and correlated with risk factors for ischemic heart disease.

3.4 Statistical Analysis

All the collected data was entered in Microsoft Excel sheet and then transferred to SPSS software ver. 17 for analysis. Qualitative data was presented as frequency and percentages. P-value < 0.05 was taken as level of significance.

4. Results

Table 1. Age group distribution amongst study population

Age group	Frequency	Percent
40 to 50 years	34	43.6
51 to 60 years	14	17.9
More than 60 years	30	38.5
Total	78	100.0

As seen in the above Table 1, most of the study population belongs to the age group of 40 to 50 years (43.6 %) followed by more than 60 years (38.5%) and 51 to 60 years (17.9%) (Age range = 40-73).

Table 2. Comorbidities amongst Study Population

Comorbidities	Frequency	Percent
Hypertension	46	59.0
Diabetes	39	50.0

As seen in the above Table 2, Hypertension was the most common clinical features amongst study population (59%) followed by Diabetes (50%).

Table 3. Habits amongst study population

Habits	Frequency	Percent
Alcohol	3	3.8
Tobacco (Mishri user)	13	16.7

As seen in the above Table 3, Habitssuch as alcohol and Tobacco chewing was observed in 3.8 % and 16.7% of study population.

Table 4. Distribution of patients according to level of Low density lipoprotein (n=78)

LDL	Frequency	Percentage
Increased (≥ 130 mg/dl)	47	60.26%
Normal (< 130 mg/dl)	31	39.74%
Total	78	100.00%

As seen in the above Table 4, Increased (≥ 130 mg/dl) LDL was observed in 60.26% of study population.

Table 5. Thyroid status findings amongst study population

Thyroid status	Frequency	Percent
Hypothyroidism	20	25.64
Hyperthyroidism	13	16.67
Normal	45	57.69
Total	78	100.00

As seen in the above Table 5, most of the study population had Normal Thyroid status (57.69%) followed by Hypothyroidism (25.64%) and Hyperthyroidism (16.67%) and SERUM TSH was the most sensitive test for evaluation.

As seen in the above Table 6, Diabetes Mellitus was statistically significant risk factor (p value < 0.05) in patients with STEMI with different levels of thyroid.

5. Discussion

Coronary artery disease is the leading cause of mortality and morbidity of both men and women accounting for over one third of total deaths²³. It has reached epidemic proportion among Indians. It accounts for 1 out of 3

women death regardless of the race or ethnicity. In women, the annual mortality rate from CAD is high²⁴.

Thyroid hormone exerts numerous effects on the cardiovascular system and is therefore not surprisingly, associated with several adverse consequences on it²⁵. Overt and subclinical hypothyroidism, with its accompanying hypercholesterolemia and hypertension, has been found to be associated with cardiovascular disease²⁶. Variations of free T3 have been linked to coronary artery disease. Several studies have shown an association between overt and subtle thyroid function abnormalities and atherosclerosis. It remains a matter of speculation that variation of thyroid function within the normal range can lead to adverse outcome with respect to Coronary Artery Disease (CAD)²⁷.

5.1 Demographic Profile

In the present study, most of the study population belongs to the age group of 40 to 50 years (43.6 %) followed by More than 60 years (38.5%) and 51 to 60 years (17.9%) with mean age of 58.12 ± 5.053 years. In the present study, most of the study population were married (91%). This findings is in agreement with the study conducted by Manish Sahni in which mean age was 62.74 ± 13.6 years and in Veena Nanjappa in which mean age was 64.4 ± 11 years^{28,29}. So as age increase risk of myocardial infarction increases in woman. This increase in incidence of IHD with age is observed in Framingham heart study.³⁰ Protection from IHD in younger women is due to endogenous estrogen. At the age of 60, the level of atherogenic lipids increase and risk of IHD doubles for women^{6,7}.

5.2 Co-morbidities

In the present study, Hypertension was the most common co-morbidities amongst study population (59%) followed by Diabetes (50%). This findings is in agreement with the study conducted by Meenakshi Rana et al., in which

Table 6. Analysis of risk factors of STEMI with different levels of thyroid

Risk Factors	Hypothyroidism		Hyperthyroidism		Normal	
	OR	P value	OR	P value	OR	P value
Diabetes	4.1	0.01	3.4	0.004	2.1	0.005
Hypertension	1.5	0.43	0.7	0.89	1.3	0.711
Dyslipidemia	0.34	0.78	0.4	0.893	1.1	0.69
Alcohol	0.6	0.69	0.3	0.632	0.7	0.87
Tobacco	0.7	0.95	0.3	0.72	0.9	0.45

diabetes Mellitus was statistically significant risk factor (p value <0.05) in patients with STEMI with different levels of thyroid (Table 6). Women with diabetes are 3–7 times more likely to develop or die of coronary heart disease than women with-out diabetes. This is much higher than the 2- to 3-fold increased risk experienced by men with diabetes. In our study diabetes is observed 0as significant risk factors in patients with STEMI with different levels of thyroid³¹.

Furthermore, Diabetes is a predictor of atypical presentation of acute myocardial infarction in women and different researches suggests that shortness of breath may be an important acute coronary syndrome symptom in women with diabetics^{32,33}. These two risk factors were found in Palestinian study in 2013 in patients with CAD that showed high prevalence of hypertension and diabetes in women 74.3%, 65.7% respectively³⁴.

Diabetes carries a greater risk in females, completely eliminating the “female advantage”³⁵. Diabetes removes the estrogens protective effects and eliminates the normal sex difference in the prevalence of IHD³⁶. Diabetes equalises the risk of IHD between premenopausal diabetic women and non-diabetic men of same age³⁷. Diabetes is associated with other IHD risk factors like obesity, dyslipidemia, hypertension, and insulin resistance. Lastly diabetes is associated with various coagulation abnormalities like endothelial dysfunction and platelet abnormalities, additional contributors to IHD³⁵.

5.3 Habits

In the present study, Habits such as alcohol and Tobacco chewing were observed in 3.8% and 16.7% of study population. Similarly, in the study conducted by Shruthi Bettegowda. in which tobacco consumption was present in 20% females either in the form of smoking or chewing.³⁸ Smoking increases the IHD risk in females by its synergistic action with oral contraceptive use, especially in women aged more than 35 years. Women with smoking attains early menopause, another IHD risk unique to females^{39,40}.

5.4 Dyslipidemia

Certain lipid components or lipoproteins may pose a greater risk in women than in men and may provide a different target for risk modification. Compared with men, Low-Density Lipoprotein (LDL) cholesterol is lower, on average, and High-Density Lipoprotein (HDL)

cholesterol is higher in premenopausal women. After menopause, LDL cholesterol concentrations rise.

5.5 Thyroid Status

In the present study, most of the study population had Normal Thyroid status (57.69%) followed by Hypothyroidism (25.64%) and Hyperthyroidism (16.67%)

In the present study, most of the study population had less than 10 days of hospital stay (78.2%) followed by more than 10 days (21.8%). Diabetes Mellitus was statistically significant risk factor (p value <0.05) in patients with STEMI with different levels of thyroid.

6. References

1. Heart Disease and Stroke Statistics-2009 Update: A Report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee.
2. Kim ESH, Carrigan TP, Menon V; Enrollment of women in National Heart, Lung, and Blood Institute-funded cardiovascular randomized controlled trials fails to meet current federal mandates for inclusion. *J Am Coll Cardiol.* 2008; 52(8):672-3. <https://doi.org/10.1016/j.jacc.2008.05.025> PMID:18702973
3. Pepine CJ. Ischemic heart disease in women. *J Am Coll Cardiol.* 2006; 47:S1-S3. <https://doi.org/10.1016/j.jacc.2005.10.022> PMID:16458166
4. Mikhail GW. Coronary heart disease in women. *BMJ.* 2005; 331(7515):467-8. <https://doi.org/10.1136/bmj.331.7515.467> PMID:16141136 PMCID:PMC1199011
5. Wenger NK. Coronary heart disease: The female heart is vulnerable. *Prog Cardiovasc Dis.* 2003; 46(3):199-299. <https://doi.org/10.1016/j.pcad.2003.08.003> PMID:14685940
6. Stampfer MJ, Colditz GA, Willett WC, Manson JE, Rosner B, Speizer FE. Postmenopausal estrogen therapy and cardiovascular disease. Ten-year follow-up from the nurses health study. *N Engl J Med.* 1991; 325:756-762. <https://doi.org/10.1056/NEJM199109123251102> PMID:1870648
7. Stampfer MJ, Colditz GA; Estrogen replacement therapy and coronary heart disease: Quantitative assessment of the epidemiologic evidence. *Prev Med.* 1991; 20(1):47-63. [https://doi.org/10.1016/0091-7435\(91\)90006-P](https://doi.org/10.1016/0091-7435(91)90006-P)
8. Enas EA, Senthilkumar A, Juturu V, Gupta R. Coronary artery disease in women. *Indian Heart J.* 2001; 53:282-92.
9. O'Toole ML. Exercise and physical activity. In Douglas PS editor. *Cardiovascular health and disease in women.* Philadelphia: W. B. Saunders; 1993.p. 253-68.
10. Manson JE, Colditz GA, Stampfer MJ, Willett WC, Rosner B, Manson RR. A prospective study of obesity and risk of coronary heart disease in women. *N Engl J*

- Med. 1990; 322(13):882-889. <https://doi.org/10.1056/NEJM199003293221303> PMID:2314422
11. Danzi S, Klein I. Thyroid disease and the cardiovascular system. *Endocrinol Metab Clin North Am.* 2014; 43(2):517-28. doi: 10.1016/j.ecl.2014.02.005. <https://doi.org/10.1016/j.ecl.2014.02.005> PMID:24891175
 12. Fazio S, Palmieri EA, Lombardi G, Biondi B. Effects of thyroid hormone on the cardiovascular system. *Recent Prog Horm Res.* 2004; 59:31-50. <https://doi.org/10.1210/rp.59.1.31> PMID:14749496
 13. Klein I, Danzi S. Thyroid disease and the heart. *Circulation.* 2007; 116(15):1725-35. <https://doi.org/10.1161/CIRCULATIONAHA.106.678326> PMID:17923583
 14. Danzi S, Klein I. Alterations in thyroid hormones that accompany cardiovascular disease. *Clinical Thyroidology.* 2009; 21(1):3-5.
 15. Boekholdt SM, Titan SM, Wiersinga WM, Chatterjee K, Basart DC, Luben R, et al. Initial thyroid status and cardiovascular risk factors: The EPIC-Norfolk prospective population study. *Clin Endocrinol (Oxf).* 2010; 72(3):404-10. <https://doi.org/10.1111/j.1365-2265.2009.03640.x> PMID:19486022
 16. Ertek S, Cicero AF. Hyperthyroidism and cardiovascular complications: A narrative review on the basis of pathophysiology. *Arch Med Sci.* 2013; 9(5):944-52. <https://doi.org/10.5114/aoms.2013.38685> PMID:24273583 PMID:PMC3832836
 17. Pimentel RC, Cardoso GP, Escosteguy CC, Abreu LM. Thyroid hormone profile in acute coronary syndromes. *Arq Bras Cardiol.* 2006; 87(6):688-94. <https://doi.org/10.1590/S0066-782X2006001900002>
 18. Coceani M. Heart disease in patients with thyroid dysfunction: hyperthyroidism, hypothyroidism and beyond. *Anadolu Kardiyol Derg.* 2013; 13(1):62-6. <https://doi.org/10.5152/akd.2013.008> PMID:23086805
 19. Grais IM, Sowers JR. Thyroid and the heart. *Am J Med.* 2014; 127(8):691-8. <https://doi.org/10.1016/j.amjmed.2014.03.009> PMID:24662620 PMID:PMC4318631
 20. Biondi B, Klein I. Hypothyroidism as a risk factor for cardiovascular disease. *Endocrine.* 2004; 24(1):1-13. <https://doi.org/10.1385/ENDO:24:1:001>
 21. Yang L, Zou J, Zhang M, Xu H, Qi W, Gao L, et al. The relationship between thyroid stimulating hormone within the reference range and coronary artery disease: Impact of age. *Endocr J.* 2013; 60(6):773-9. <https://doi.org/10.1507/endocrj.EJ12-0448> PMID:23470525
 22. Thom T, Hasse N, Rosamond W. Heart disease and stroke statistics-2006 update: A report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation.* 2006; 113:e85. <https://doi.org/10.1161/CIRCULATIONAHA.105.171600> PMID:16407573
 23. Kretsoulas Catherine, Sloane Debi, Pogue Janic, Velianou James L., Anand Sonia S. Referrals in acute coronary events for cardiac catheterization: The RACE CAR Trial. *Can J Cardiol.* 2010; 8:e290-e296. [https://doi.org/10.1016/S0828-282X\(10\)70436-0](https://doi.org/10.1016/S0828-282X(10)70436-0)
 24. Thom T, Hasse N, Rosamond W. Heart disease and stroke statistics-2006 update: A report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation.* 2006; 113:e85. <https://doi.org/10.1161/CIRCULATIONAHA.105.171600> PMID:16407573
 25. Braverman LE, Cooper D. 7th ed. Wolters Kluwer Health. Werner & Ingbar's *The Thyroid: A Fundamental and Clinical Text*; 2012.
 26. Cappola AR, Ladenson PW. Hypothyroidism and atherosclerosis. *J Clin Endocrinol Metab.* 2003; 88(6):2438-44. <https://doi.org/10.1210/jc.2003-030398> PMID:12788839
 27. Taylor PN, Razvi S, Pearce SH, Dayan CM. Clinical review: A review of the clinical consequences of variation in thyroid function within the reference range. *J Clin Endocrinol Metab.* 2013; 98:3562-3571 <https://doi.org/10.1210/jc.2013-1315> PMID:23824418
 28. Sahni M, Kumar R, Thakur S, Bhardwaj R. Clinical profile, risk factors and shortterm outcome of acute myocardial infarction in females: A hospital based study. *Heart India.* 2013; 173-7
 29. Nanjappa V et al. Clinical profile and 30-day outcome of women with acute coronary syndrome as a first manifestation of ischemic heart disease: A single-center observational study. *Indian Heart J.* 2016 Mar-Apr; 68(2):164-8. <https://doi.org/10.1016/j.ihj.2015.08.006> PMID:27133325 PMID:PMC4867941
 30. Castelli WP, Kannel WB. Cardiovascular disease in women. *Am J Obstet Gynecol.* 1988; 138:1553. [https://doi.org/10.1016/0002-9378\(88\)90189-5](https://doi.org/10.1016/0002-9378(88)90189-5)
 31. Rana M, Pandey M, Devpura G. Study of risk factors for st segment elevation myocardial infarction and it's correlation with thyroid functions profile. *Ann Int Med Den Res.* 2018; 4(2):ME53-55. <https://doi.org/10.21276/aimdr.2018.4.2.ME13>
 32. Stephen SA, Darney BG, Rosenfeld AG. Symptoms of Acute Coronary Syndrome in Women with Diabetes: An Integrative Review of the Literature. *Heart Lung.* 2008; 37:179-89. <http://dx.doi.org/10.1016/j.hrtlng.2007.05.006> PMID:18482629
 33. Culic V, Eterovic D, Miric D, Silic N. Symptom presentation of acute myocardial infarction: influence of sex, age, and risk factors. *American Heart Journal.* 2002; 144:1012-7. <https://doi.org/10.1067/mhj.2002.125625> PMID:12486425

34. Jamee A, Abed Y, Jalambo MO. Gender difference and characteristics attributed to coronary artery disease in gaza-palestine. *Global Journal of Health Science*. 2013; 5:51-6. <https://doi.org/10.5539/gjhs.v5n5p51> PMID:23985106 PMCID:PMC4776843
35. Jousilahti P, Vartiainen E, Tiromilehto J, Puska P. Sex, age, cardiovascular risk factors, and coronary heart disease: A prospective follow up study of 14,786 middle aged men and women in Finland. *Circulation*. 1999; 99(9):1165-72. <https://doi.org/10.1161/01.CIR.99.9.1165> PMID:10069784
36. Sowers JR. Diabetes mellitus and cardiovascular disease in women. *Arch Intern Med*. 1998; 158:617-621. <https://doi.org/10.1001/archinte.158.6.617> PMID:9521226
37. Dhar M, Dwivedi S, Agarwal MP, Rajpal S. Clinical profile of coronary artery disease in women. *Indian J Cardiology*. 2006; 9:18-23.
38. Bettgowda S. Clinical profile of ischemic heart disease in women with special reference to the risk factors sch. *J App Med Sci*. 2014; 2(6C):3020-5.
39. Colditz GA, Bonita R, Stamfer MJ, Willet WC, Rosner B, Speizer FE, Hennekens CH. Cigarette smoking and risk of stroke in middle aged women. *N Engl J Med*. 1988; 318(15):937-41. <https://doi.org/10.1056/NEJM198804143181501> PMID:3352685
40. Kaul U, Dogra B, Manchanda SC, Wasir HS, Rajani M, Bhatia ML. Myocardial infarction in young Indian patients: Risk factors and coronary arteriographic profile. *Am Heart J*. 1986; 112(1):75-75. [https://doi.org/10.1016/0002-8703\(86\)90680-0](https://doi.org/10.1016/0002-8703(86)90680-0)

How to cite this article: Chafekar, N., Babras, M. and Kirloskar, M. Study of the Risk Factors for Coronary Artery Disease and their Correlation with Thyroid Hormone Profile amongst Women with ST Segment Elevation in Acute Myocardial Infarction. *MVP J. Med. Sci.* 2021; 8(1): 140-146.